

# Scientific Consensus Statement on Environmental Agents Associated with Neurodevelopmental Disorders



*Developed by the Collaborative on Health and the Environment's  
Learning and Developmental Disabilities Initiative*

*February 20, 2008  
(revised July 1, 2008)*

*Given established scientific knowledge, protecting children from neurotoxic exposures from the earliest stages of fetal development is clearly an essential public health measure. By reducing environmental factors that may lead to learning and developmental disorders, we will create a healthier environment in which all children can reach and maintain their full potential.*

**Drafted and edited by:**

Steven G. Gilbert, PhD, DABT, Institute of Neurotoxicology and Neurological Disorders

**Scientific Review Committee:**

David C. Bellinger, PhD, MSc, Harvard Medical School  
Lynn R. Goldman, MD, MPH, Johns Hopkins University, Bloomberg School of Public Health  
Philippe Grandjean, MD, Harvard School of Public Health  
Martha Reed Herbert, MD, PhD, Massachusetts General Hospital  
Philip J. Landrigan, MD, MSc, Mount Sinai School of Medicine  
Bruce P. Lanphear, MD, MPH, University of Cincinnati  
Barbara McElgunn, RN, Learning Disabilities Association of Canada  
John Peterson Myers, PhD, Environmental Health Sciences  
Isaac Pessah, PhD, University of California, Davis  
Ted Schettler, MD, MPH, Science and Environmental Health Network  
Bernard Weiss, PhD, University of Rochester School of Medicine and Dentistry

**Other Reviewers:**

Aimee Boulanger, Institute for Children's Environmental Health  
Sean Foley, Toxipedia, Institute of Neurotoxicology and Neurological Disorders  
Elise Miller, MEd, Institute for Children's Environmental Health and the Collaborative on  
Health and the Environment's Learning and Developmental Disabilities Initiative  
Nancy Snow, MS, Institute for Children's Environmental Health

*Note: The reviewers listed above provided comments and suggestions during the drafting process, but do not necessarily agree with every aspect of this paper. The signatories at the end of the paper are those who fully agree with and support this statement.*

## Contents

1 Introduction.....	3	4.2.11 Nicotine and environmental tobacco smoke.....	13
1.1 Purposes of the document.....	3	4.3 Other contributors and emerging evidence.....	13
1.2 Scope and incidence of disorders.....	3	4.3.1 Endocrine disruptors.....	13
1.2.1 Definitions of LDDs.....	4	4.3.2 Fluoride.....	14
1.2.2 Overview of brain development.....	4	4.3.3 Food additives.....	14
1.2.3 Incidence.....	4	5 Call for Further Research.....	15
1.2.4 Impacts on individuals, families and society.....	5	5.1 Better assessment tools and procedures..	15
2 Historical Perspective.....	5	5.2 In vitro (test-tube) screening of both new and old compounds.....	15
3 Environmental Contributors to Disorders.....	7	5.3 Chronic effects of hazardous chemicals..	15
3.1 Timing of exposure.....	7	5.4 Multiple exposures.....	15
3.2 Bioaccumulation and mixtures.....	7	5.5 Low-level exposures.....	15
3.3 Mechanisms of disruption.....	8	5.6 Interactions with socioeconomic factors (environmental justice).....	16
3.4 Variable sensitivity.....	8	5.7 Effects of endocrine disruption on cognitive deficits.....	16
3.5 Evidence of exposures (body burden).....	8	5.8 Interactions with genetics and identification of susceptible subpopulations..	16
3.6 Other environmental factors.....	8	5.9 World Health Organization research recommendations.....	16
3.6.1 Pharmaceuticals.....	9	6 Ethical and Policy Considerations.....	16
3.6.2 Genetic factors.....	9	6.1 Ethical considerations.....	16
3.6.3 Environmental justice: socioeconomics, nutrition and stress.....	9	6.2 Policy considerations.....	16
4 Status of Evidence on Environmental Agents..	9	7 Conclusions.....	17
4.1 Human and animal assessment of learning and development.....	9	8 Resources: Children and Environmental Agents.....	18
4.2 High-confidence conclusions.....	10	9 Glossary of Terms and Term Usage Specific to This Statement.....	19
4.2.1 Alcohol.....	11	10 Acknowledgments.....	23
4.2.2 Lead.....	11	11 References.....	24
4.2.3 Mercury.....	11		
4.2.4 PCBs.....	12		
4.2.5 PBDEs.....	12		
4.2.6 Manganese.....	12		
4.2.7 Arsenic.....	12		
4.2.8 Solvents.....	12		
4.2.9 PAHs.....	13		
4.2.10 Pesticides.....	13		

# 1 Introduction

This consensus statement outlines the current scientific understanding of the links between environmental factors and learning and developmental disabilities. It also identifies important research areas that hold promise of further advancing our understanding of these links. This statement is intended as a guide to scientists, medical professionals, policymakers, public health advocates, and the general public in advancing their efforts to address the important individual and social issues raised by learning and developmental disabilities.

Terms in orange underlined font are defined in a Glossary of Terms and Term Usage specific to this document, beginning on page 19.

## 1.1 Purposes of the document

- To review findings from diverse research disciplines concerning environmental contaminants and the biological basis of compromised learning and development, with special attention to critical recent discoveries in related basic sciences;
- To identify conclusions that could be drawn with confidence from existing data;
- To identify critical knowledge gaps and areas of uncertainty;
- To establish key elements of a coherent research agenda to help fill these gaps and resolve uncertainties;
- To form a foundation of current scientific knowledge upon which to make policy decisions that promote and protect an environment in which all children can reach and maintain their full potential.

## 1.2 Scope and incidence of disorders

Within the human brain lies our capacity to learn, talk, read, calculate, memorize, conceptualize, organize, pay attention, utilize motor skills, interact socially and behave appropriately. We cannot reach our full potential with a damaged brain or nervous system.

Though many of the environmental contaminants discussed in this document can also undermine the healthy development of other biological systems in the body, such as the reproductive, endocrine and immune systems, this consensus statement is focused solely on the developing brain and nervous system. This distinction is somewhat artificial since the impact of toxicants upon these other systemic processes may have effects that feed back and impact brain function. Although it is often believed that endocrine function, for example, is independent of brain development, it is now clear that they are so closely entwined that endocrine toxicants are, in essence, neurodevelopmental toxicants.

### Children are not little adults



- Environmental exposures start early: pre-conception, during gestation (*in utero* exposure), via breast milk, infant formula and then through contact with the environment.
- For their body weight, children eat and breathe more than adults, thus a small exposure translates into a big dose.
- Their organ systems, particularly the nervous system, are forming and are thus more susceptible to the effects of chemicals.
- Young children are prone to hand-to-mouth behaviors that expose them to higher levels of ambient chemicals.
- Children rely on adults to ensure that they develop in an environment in which they can reach and maintain their full potential.

Other neurologically based disorders have also been shown to have environmental contributors, especially Parkinson's disease (1-4), mental illness including schizophrenia (5), cerebral palsy (6) and epilepsy. This document focuses on learning and developmental disorders and does not address these conditions.

### 1.2.1 Definitions of LDDs

In this report, we define learning and developmental disorders broadly as conditions resulting from interference of normal brain development and function that adversely affect an individual's performance. [Learning and developmental disabilities \(LDDs\)](#) include but are not limited to deficits in learning and memory, reduced IQ, [attention deficit hyperactivity disorder \(ADHD\)](#), [autism spectrum disorder](#), [conduct disorders](#) and [developmental delays](#).

### 1.2.2 Overview of brain development

The development of the human brain begins in utero and continues through adolescence, following a precise and delicate step-by-step sequence involving complex neurobiological processes including the formation of the neural tube, cell proliferation, differentiation, migration and selection, synapse formation, development of neurochemical systems, cell pruning; and myelination. These processes ultimately involve 10 to 100 billion neuronal cells with many trillions of connections. The long and complex development of the brain and nervous system leaves it susceptible to the adverse effects of chemical exposures.

Even minor changes in the structure or function of the nervous system can have profound consequences for neurological, behavioral and related body functions. Disruption of the brain's normal development can happen in utero as a baby develops within the mother's body or as the brain continues to develop from infancy through adolescence. Brain function can also be impacted in adulthood from an accident, trauma or exposure to [neurotoxicants](#).

### 1.2.3 Incidence

[Autism](#), [attention deficit hyperactivity disorder \(ADHD\)](#), [dyslexia](#), [mental retardation](#), lowered IQ and other disorders of learning and behavior are highly prevalent among American children. The incidence of [learning and developmental disabilities \(LDDs\)](#) appears to be rising, affecting between five and 15 percent of all children under the age of 18 in the United States, or more than 12 million children under 18 (7). In general, disabilities have increased significantly over the past four decades (8).

These disabilities include disorders of learning, attention, emotional state and behavior. [Mental retardation](#) impacts two percent of children, or approximately 1.4 million children (9-12). [ADHD](#) is conservatively estimated to occur in three to six percent, or approximately two million children (13) (14). A more recent study in the [National Health and Nutrition Examination Survey \(NHANES\)](#) found 8.7 percent of 8- to 15-year-old children met the Diagnostic and Statistical Manual of Mental Disorders ([DSM-IV-TR](#)) criteria for ADHD based on parent recall (15).


#### Critical recent discoveries

- Even very low doses of some biologically active contaminants can alter gene expression important to learning and developmental function.
- Exposures during fetal development can adversely affect learning and development of the individual and last a lifetime.
- Humans are exposed to complex mixtures of chemicals that can interact to enhance adverse effects.
- Due to genetic variation ([polymorphisms](#)) people differ in susceptibility to exposures. Not identifying and studying susceptible subgroups can result in failure to protect those at high risk.
- Children are often more susceptible than adults to the effects of exposure to [environmental agents](#).

Autism spectrum disorder is estimated to affect approximately 0.7 percent, or 450,000 children and appears to be 10 times more prevalent today than it was in the 1980s (16), although that estimate is still debated.

These nonfatal disabilities that affect an individual for a lifetime can be classified as the “new morbidities,” and the prevalence in children is summarized in Figure 1 below. This figure presents a compilation of data from different sources (7, 15, 17-19), demonstrating the interrelation of these childhood conditions.

### Prevalence of the “New Morbidities” in United States Children



#### 1.2.4 Impacts on individuals, families and society

In children, developmental, learning, attention and behavioral problems can cause tremendous challenges for the affected children, their families and communities. Consequences include psychological and economic costs associated with learning delays, aggressive or otherwise inappropriate behavior, school dropout, teen parenting, substance abuse, unemployment, welfare dependency and involvement with juvenile and adult criminal justice systems (20).

Attempts to calculate the costs of these childhood “morbidities” have only recently been undertaken (20, 21). Providing special education services to students with disabilities amounted to \$77.3 billion, or an average of \$12,474 per student in 1999-2000, almost twice the cost per regular education student, which is almost 22 percent of the 1999-2000 total spending on all elementary and secondary educational services in the US (22) (for additional information see the Center for Special Education Finance (CSEF) at <http://www.csef-air.org/>).


## 2 Historical Perspective

The effects of chemicals on the nervous system have been recognized since ancient times. Most of the early reports were the results of high exposures in workers that led to debilitating conditions or even death. For example, 2000 years ago people knew that lead exposure “makes the mind give way.” As

science progressed, it was recognized that even small doses of some chemicals result in subtle nervous system impacts that affect an individual's performance. Despite this knowledge, lead was added to paint and gasoline, removed only following considerable research that confirmed what was already known. Many investigations have now demonstrated that even small amounts of lead affect the developing nervous system (23).

Similarly, the adverse effects of exposure to elemental mercury, recognizable because of its liquid silver appearance, which evaporates at room temperature and can be inhaled, were recognized in miners well before the common era. Subsequently, workers in the felt-hat industry, where mercury was used to process the fur, succumbed to its **neurotoxic** effects – truly an example of “mad hatters” (see *Alice's Adventures in Wonderland* by Lewis Carroll). Organic mercury, formed by the conversion of inorganic mercury to methylmercury by bacteria, **bioaccumulates** in fish and is readily absorbed. Organic mercury exposures in Iraq in 1971 and in Minimata, Japan, beginning in the late 1950s revealed both the overt and more subtle effects of organic mercury on development and the nervous system. Even low doses of methylmercury damage the nervous system and cause neurobehavioral deficits (24-28).

Exposure to a wide range of potentially hazardous chemicals from conception to death is now unavoidable. Approximately 3,000 chemicals are produced in quantities greater than one million pounds per year. For the majority of these chemicals little information exists about the potential effects on learning and development. There is good evidence that about 200 of these chemicals are adult neurotoxicants and another 1000 are suspected of affecting the nervous system (29) (see Figure 2, in part from (29)). Overall there has been a gross failure to require developmental neurotoxicity testing.



As our testing methods have become more sophisticated, the recognition of individual sensitivity and, in particular, the sensitivity of the developing nervous system to the effects of [environmental agents](#) has grown. The causes of and initiatives to prevent [LDDs](#) have received increased attention from national organizations (30-32), the government and academic researchers (20, 29, 33). A substantial recent initiative, the [National Children's Study](#), will examine the effects of environmental influences on the health and development of more than 100,000 children across the United States, following them from before birth until age 21, with the goal of improving the health and well-being of children (<http://www.nationalchildrensstudy.gov/>).

### **3 Environmental Contributors to Disorders**

Many factors – heredity, gene expression, socioeconomic environment, infectious disease, nutrition, stress, drugs and chemical contaminants – contribute in complex ways to brain development and thus to the genesis of [LDDs](#). There is growing evidence that the interaction of these factors is associated with or exacerbates a variety of developmental disorders (34-37). Of all the factors contributing to [LDDs](#), chemical contaminants in the environment have historically been the least studied despite being the most preventable. We now have solid scientific evidence that a variety of [environmental agents](#) can adversely affect the nervous system. The impact of chemical contaminants on children is a worldwide issue and the focus of this document.

The susceptibility of the developing nervous system to the adverse effects of environmental agents, and the need to take preventive measures were recently recognized by international researchers in the Faroes Statement (38). International groups have also developed statements on the developmental effects of metals and the need to take preventive action to protect public health (39). In addition, comprehensive reports have documented the effects of environmental agents on development (40).

#### **3.1 Timing of exposure**

A child's developing nervous system is more sensitive to chemical exposures than the adult nervous system. This can be seen in the effects of alcohol: a pregnant woman who drinks enough to become intoxicated may suffer a hangover, but her fetus may suffer permanent brain damage resulting in fetal alcohol spectrum disorder (41). Also, in both the Iraq and Minimata methylmercury disasters, pregnant women showed minimal signs of toxicity, but their children displayed effects ranging from cerebral palsy to delayed development (6). Extensive animal data also establish the effects of mercury exposure on the developing nervous system.

Research on brain development has mapped out the progression of the proliferation, differentiation and migration of different cell types into selected regions of the brain. This work has been extended as neurotoxicologists have examined the effects of exposure to chemicals at different points during development (42-45). Vulnerability to chemical exposure varies across the stages of brain development (43) such that exposure at three months gestation may result in a different effect than exposure to the same chemical at six months gestation or at two years of age.

#### **3.2 Bioaccumulation and mixtures**

Concentrations and/or potency of [environmental agents](#) can be amplified because of [persistence](#) ([biomagnification](#) and [bioaccumulation](#)) and because agents occur in [mixtures](#). Mercury accumulation in fish is a well known example of both [biomagnification](#) and [bioaccumulation](#) (24, 46). Fish can also be contaminated with other compounds, such as [polychlorinated biphenyls \(PCBs\)](#); thus we are often consuming a [mixture](#) of compounds. Recent [biomonitoring](#) studies reveal the range of compounds we are exposed to and that accumulate in our bodies. Experiments with single chemicals can underestimate the effects of these chemicals in [mixtures](#).

The interplay among multiple toxicants whether stored or through ongoing exposures, and other environmental factors can cumulatively interfere with the brain's development and exacerbate the impact. For example, there is good evidence that lead exposure and maternal stress interact, and there is some evidence of [potentiated](#) effects of combined exposures of the pesticides paraquat and maneb during development (47).

### 3.3 Mechanisms of disruption

Brain development is a long and complex process that involves cellular proliferation, migration, differentiation, synaptogenesis, myelination and apoptosis (programmed cell death). Chemicals can interfere with or stop these processes through the extended period of brain development from conception to adolescence (43). There are multiple mechanisms by which chemicals can disrupt neurological development, including gene expression acting on protein pathways (48) (49, 50) and hypothyroidism (51-53). Recent research has focused on how genes act on protein pathways, determining how tissue and cells form and grow. The expression of multiple genes can affect multiple protein pathways. A single contaminant that influences gene expression can affect [endpoints](#) in more than one tissue. Some contaminants have been shown to alter the expression of hundreds of genes, and effects can vary with the timing and dose of the contaminant.

### 3.4 Variable sensitivity

Genetic variation, or DNA [polymorphisms](#), within populations (humans, wildlife and laboratory animals) can result in greater sensitivity to specific contaminants in some individuals. Specific [genetic polymorphisms](#) are linked to increased risk to various disorders such as fetal alcohol spectrum disorder (54) and are associated with increased susceptibility to organophosphate pesticide toxicity (55-57). Genetic variations are also associated with increased susceptibility to higher blood-lead levels (58, 59).

The effects of environmental agents, particularly endocrine disruptors, have recently been shown to promote epigenetic transgenerational effects – changes passed down to subsequent generations. [Epigenetics](#), a molecular phenomenon, typically methylation of the genome, regulates gene expression without alterations to the DNA sequence (60, 61). Recent studies have demonstrated that the commonly used fungicide vinclozolin has produced epigenetic transgenerational effects on development and cancer (62-64). Epigenetic effects may explain in part the transgenerational effects of the synthetic estrogen diethylstilbestrol (65). The low-dose effects of endocrine disruptors are being actively explored (66, 67).

### 3.5 Evidence of exposures (body burden)

Recent measurements of contaminants in people demonstrate that humans are exposed, starting at conception, to hundreds of chemicals simultaneously – and some at levels within ranges known individually to affect neurological development (68-70).

Chemicals may also be passed from one generation to the next, from mother to developing child, as chemicals stored in or consumed by the mother cross the placenta or are incorporated into breast milk. While breast milk is clearly the best food for infants and breast feeding is almost always recommended, [persistent, bioaccumulative](#) chemicals stored in a woman's body fat are mobilized during breast feeding and excreted in breast milk.

### 3.6 Other environmental factors

There are many other chemicals of concern, as well as other environmental factors that interact with chemicals, that can affect development and contribute to LDDs. We include a brief discussion of some of

these factors due to increasing evidence of their interactions with chemical agents and their contribution to LDDs.

### **3.6.1 Pharmaceuticals**

A wide range of drugs have [neurotoxic](#) side effects in children when consumed by a pregnant mother, infant or child. When these drugs are given to children (even when needed) and not adequately monitored, serious lifetime disabilities can result. A classic example is aminoglycoside antibiotics, which can cause severe hearing loss and subsequent learning and development challenges. We know much less about the long-term effects of psychoactive compounds. There is evidence that thalidomide or valproate exposure during pregnancy increases the risk of autism (71-74).

### **3.6.2 Genetic factors**

With [autism](#) in particular, new research is beginning to reveal that the disorder involves the whole body, a complex interaction of genes and the environment, and perhaps many factors working in concert with one another (75, 76). While the low-dose effects of endocrine disruptors are still being explored, some have argued that genetic susceptibility plays an important role in the etiology of autism (76).

### **3.6.3 Environmental justice: socioeconomics, nutrition and stress**

Six million children live in poverty in the United States, increasing the likelihood of exposures and heightened vulnerability to [environmental agents](#) that adversely affect learning and development. Economically disadvantaged children are more likely to live in older homes contaminated with lead, live in neighborhoods and around schools where pesticides have traditionally been applied, and have diets that are less nutritious. Additionally, there is growing evidence that stress combined with environmental exposures increases susceptibility to developmental disorders (36, 47). Recent research indicates that stress and social ecology can play an important role in developmental disorders (36).

Children lacking certain nutrients are more vulnerable to toxicants. For example, iron and/or calcium deficiency affects the absorption and toxicity of heavy metals such as lead (77) and manganese (78, 79). Lead is stored in bone and may be mobilized with calcium from bone to the fetus during pregnancy. The role of nutrition in mitigating exposure to environmental agents is an important public health issue (80).

## **4 Status of Evidence on Environmental Agents**

Research definitively shows that [environmental agents](#) such as lead, mercury, manganese, arsenic, [PCBs](#), alcohol, toluene, tobacco smoke and many pesticides are capable of disrupting human brain development, resulting in negative impacts on the functions controlled by the brain. Additional environmental chemicals and pollutants, other solvents and other heavy metals have been shown to disrupt brain development in animal studies and are suspected of having similar effects in humans.

### **4.1 Human and animal assessment of learning and development**

Evaluating the potential [neurotoxic](#) effects of a compound often requires evaluating animal data and then incorporating any human data into the assessment.

Given the large number of compounds with limited or no data on nervous system effects, it is important to consider [in vitro](#) testing. [In vitro](#) systems typically use cell culture techniques for initial assessment of [neurotoxic](#) potential. New tests utilizing cell lines are now used to assess the effects of chemicals on eye irritation or damage. Cell lines are available with which to examine a variety of [endpoints](#) on neuron or glial cells. [Endpoints](#) include proliferation, migration, synaptogenesis and apoptosis. Well-characterized [in vitro](#) testing systems have a number of potential advantages including minimizing the use of animals, reducing costs and increasing adaptability to rapidly screen agents for potential cellular effects. In vitro

tests are not sufficient for setting exposure standards, but they can provide a rapid assessment that further testing is needed.

It is also possible to use [in vitro](#) models to examine chemical [mixtures](#), such as additives, on neuronal cells (81). There are, however, a number of challenges with [in vitro neurotoxicity](#) testing, such as characterizing cell migration and interconnection (82).

Several different tests are used to assess [neurotoxic](#) effects in adult and developing animals. Functional assessment can be supplemented with morphological assessment of the nervous system (83) although conventional pathological assessments are relatively insensitive and may not detect subtle adverse cellular changes. A basic screen for behavioral function and neurological involvement is the functional observational battery (FOB) (84). The FOB is typically used to evaluate the need for more sophisticated [neurotoxicity](#) testing.

The US Environmental Protection Agency (EPA) and the Paris-based Organisation for Economic Co-operation and Development (OECD) established a protocol for the evaluation of developmental [neurotoxicity](#) (DNT) in laboratory animals (US EPA 870.6300 and OECD 426) (85, 86). These protocols include tests of neurobehavioral function, auditory startle (hearing), learning and memory function, changes in motor activity, [neuropathologic](#) examination and [morphometric](#) analysis. For examples and comments on the testing protocols see papers by Claudio et al., Cory-Slechta et al, Dorman et al., Garman et al. and Mileson and Ferenc (87-91).

Testing involving animals, including nonhuman primates, has been invaluable in evaluating the effects of a number of [neurotoxicants](#) (92). Recent studies have examined the [neurotoxicity](#) of pesticide [mixtures](#) in animals (93), and protocols are available to examine cognitive effects on weanling rodents (94). Advanced assessment of learning and memory in rodents has been used to evaluate the effects of lead (95). Tests of specific functions using nonhuman primates are used to evaluate the low-level effects of [neurotoxicants](#), such as mercury, on vision, auditory function and vibration sensitivity (96-99) or lead on learning and memory (100, 101). The concordance between human and animal [neurotoxicity](#) assessment is remarkable as demonstrated for lead, mercury and [PCBs](#) (102, 103).

Despite significant challenges, human testing for the [neurotoxic](#) effects of occupational exposures to chemicals is steadily advancing (104-107). These procedures have also been used to examine the [neurotoxic](#) effects of the stress and hazards of war (108, 109), which demonstrates broadening utility and acceptance of these testing procedures. The World Health Organization (WHO) recommended a neurobehavioral test battery for humans (110) and test batteries for assessing children (111). Notable examples of childhood [neurotoxic](#) evaluation include examination of the effects of low-level lead exposure (112, 113) and mercury exposure (114).

## 4.2 High-confidence conclusions

Many environmental contaminants have been conclusively shown to affect the developing nervous system, causing a range of performance deficits (29, 115).

### **Environmental agents that we are confident cause learning and developmental disabilities in humans**

- **Alcohol**
- **Lead**
- **Mercury**
- **PCBs**
- **PBDEs**
- **Manganese**
- **Arsenic**
- **Solvents**
- **PAHs**
- **Pesticides**
- **Nicotine & ETS**

### 4.2.1 Alcohol

The effects of ethyl alcohol on brain development and function are well established. Fetal Alcohol Syndrome (FAS), now considered part of Fetal Alcohol Spectrum Disorder (FASD), is the most preventable form of behavioral and learning disabilities. In the US, FASD is estimated to affect 9.1 per 1000 infants (41), with even higher rates in other parts of the world (116). Even low or moderate consumption of alcohol during pregnancy can cause subtle and permanent performance deficits (117, 118). Specific [genetic polymorphisms](#) enhance the risk of FASD (54).

### 4.2.2 Lead

Lead is probably the most studied of environmental contaminants in both humans and animals. Its effects on learning and development are undisputed. Recent research indicates that there is no safe level of lead exposure for children (112, 119, 120). Lead exposure impairs overall intelligence as measured by IQ, learning and memory and is associated with [ADHD](#) even at minute exposures. Efforts to prevent lead exposure provide an outstanding example of the struggle when science meets policy (23, 121, 122). Figure 3




Figure 3

(right) documents the gradual recognition of the harmful effects of lead at ever lower exposures. This debate is still unfolding: the US Centers for Disease Control and Prevention (CDC) has not adjusted the blood-lead action level since 1990 despite scientific evidence of behavioral effects well below 10 micrograms per deciliter ( $\mu\text{g}/\text{dL}$ ). Arguments have been made to reduce the CDC blood-lead action level to 2  $\mu\text{g}/\text{dL}$  (23).

### 4.2.3 Mercury

There is no doubt that mercury exposure causes learning and developmental disorders; the controversy regards the level of exposure. We are all exposed to some form of mercury. Inorganic mercury is the liquid silver form and is used in dental amalgams. Mercury is also present in coal, and coal-burning electric utilities facilities are a significant source of atmospheric environmental mercury. While much of the mercury falls close to the facility, mercury can be carried long distances to pollute water supplies and, ultimately, contaminate the food supply. Inorganic mercury is converted to the organic methylmercury and [bioaccumulates](#) in the flesh of fish, being [biomagnified](#) up the food chain. Methylmercury contamination often results in fish-consumption advisories, particularly for women and children. The knowledge (and concern) that methylmercury exposures affect the developing nervous system resulted in several very sophisticated studies designed to assess the effects of very low-level exposures on a range of learning and memory tests and on other performance-based tests (6, 27, 28, 46, 102, 103, 123-127). These tests typically included age-related assessment of learning and memory, reading, IQ and other neurological functions.

**4.2.4 PCBs**

Polychlorinated biphenyls (PCBs) are mixtures of chlorinated compounds that were once used as cooling and insulating fluids in electrical transformers and other electronic components. Because they are very persistent, PCBs have become widely distributed in the environment despite being banned in the 1970s. Because PCBs bioaccumulate in fat, human exposure continues through the food supply, and infant exposure continues through contaminated breast milk. Numerous studies have documented that PCB exposure can adversely affect motor skills, learning and memory as shown in lower full-scale and verbal IQ scores and reading ability (103, 128-134).

**4.2.5 PBDEs**

Polybrominated diphenyl ethers (PBDEs) have been used commonly as flame-retardant chemicals for several decades. PBDEs, structurally similar to PCBs, bioaccumulate in animals and humans, and are excreted in human breast milk. Recent studies have left little doubt that PBDEs are developmental neurotoxicants in animals and lead to changes in motor activity and reduced performance on learning and memory tests (135, 136).

**4.2.6 Manganese**

Manganese is a trace element which is necessary in small amounts for growth and development. Recent studies indicate that high levels of manganese exposure, either from inhalation (welding fumes) (137, 138) or through drinking water, can damage the developing nervous system (139, 140) as measured in full-scale IQ and verbal tests (139, 141-146). For example, a case study documented memory effects in a child exposed to manganese in drinking water (142), and a more recent study confirmed similar effects (140). The US EPA advises that water levels of manganese should not exceed 300 µg Mn/L, but approximately six percent of domestic household wells exceed this level (146).

**4.2.7 Arsenic**

Arsenic is commonly found in drinking water around the world, sometimes in concentrations high enough to cause cancer (147). Recent studies have found a dose-response relationship between exposure to arsenic and intellectual impairment (141, 148-151). While additional studies assessing the impact of low levels of arsenic in drinking water are needed, it is clear that arsenic affects the neurodevelopment of children.

**4.2.8 Solvents**

Solvents include a broad array of different compounds including toluene, benzene, alcohol, turpentine, acetone and tetrachloroethylene (TCE) (see Table 1), with more than 50 million metric tons used in the US and more than 10 million people exposed in the workplace. Solvent neurotoxicity is well recognized in adult workers (152). Ethyl alcohol is a widely used and consumed solvent with clear learning and developmental effects (see above). Recent studies indicate that occupational exposure to solvents in salons and laboratories can result in visual deficits in offspring (153-156). Several reports have documented that the adverse developmental effects

Products that are mostly solvent	Partially solvent-based
Gasoline	Glues
Diesel fuel	Adhesives
Charcoal lighter fluid	Oil-based paints
Lantern fuel	Fingernail polish
Grease	Furniture polishes
Lubricating oils	Floor polishes and waxes
Degreasing agents	Spot removers
Paint strippers	Metal and wood cleaners
Paint thinner	Correction fluid
Turpentine	Computer disk cleaners
Nail polish remover	Varnishes and shellacs
Rubbing alcohol	Wood and concrete stains

Table 1. Examples of Solvents

of maternal toluene exposure include low birth weight, decreased head circumference and [developmental delays](#) (157, 158). Awareness of developmental effects of solvent exposure has resulted in increasing concern for women working in nail and beauty salons. Some solvents, such as toluene, have also been abused by pregnant women who purposely sniff them.

#### **4.2.9 PAHs**

Polycyclic aromatic hydrocarbons (PAHs) are widely distributed air pollutants and well-recognized human mutagens and carcinogens. PAHs are generated during combustion of fuels from motor vehicles, coal-fired power plants, residential heating and cooking and are also present in tobacco smoke. Recent studies have indicated that elevated exposure to PAHs results in lower birth weight (159) and affects cognitive development (160).

#### **4.2.10 Pesticides**

Major classes of pesticides are specifically designed to kill insects, plants, fungi or animals. Agricultural and residential application of pesticides in the US totals more than one billion pounds per year, with thousands of people exposed every year. Data from acute exposure incidents leave no doubt that some pesticides, particularly insecticides, are [neurotoxic](#). There is now evidence that childhood exposure to pesticides, such as organophosphates, enhances the risk for developmental disorders including deficits in memory (161), poorer motor performance (111, 162) and an array of other conditions (163-171). A recent study documented the developmental effects of the pesticide chlorpyrifos on inner-city children (172). There is also evidence of specific genetic susceptibility to pesticide exposure and related health effects (55-57, 163).

#### **4.2.11 Nicotine and environmental tobacco smoke**

Many studies link maternal smoking during pregnancy to behavioral disorders in children (37, 173-175), and [developmental delays](#) caused by environmental tobacco smoke (ETS, also known as secondhand smoke) are costly and preventable (176). Furthermore, new data indicates that childhood exposure to ETS is associated with neurobehavioral effects (177). There is growing recognition of subsequent behavioral disorders in young adults following exposures either prenatally or as children. (175, 178, 179). The CDC reported in 2002 that 11.4 percent of all women giving birth in the United States smoked during pregnancy (180). Clearly this highly preventable form of developmental disorder requires that parents, both male and female, be educated about the harmful effects of tobacco.

### **4.3 Other contributors and emerging evidence**

It is not possible to address all the chemicals that might be associated with causing learning and developmental disorders. A more comprehensive assessment of developmental [neurotoxicity](#) of chemicals was undertaken by Grandjean and Landrigan (29) in which they pointed out that, for the majority of chemicals, we do not have the data necessary to conclude there are no adverse developmental effects. They estimate that more than 200 chemicals are known to cause neurotoxic effects in adults and that, for many of these chemicals, developmental effects have not been examined. In addition, very few studies have focused on the potential synergistic impacts of chemicals in [mixtures](#). Highlighted below are just a few agents that are of significant concern.

#### **4.3.1 Endocrine disruptors**

Animal studies have documented that a wide range of chemicals have the ability to disrupt endocrine function in animals and affect cognitive function (50). [Endocrine disruptors](#) include [phthalates](#), [PCBs](#) and polychlorinated dibenzodioxins, brominated flame retardants, [dioxins](#), [DDT](#), [perfluorinated compounds \(PFCs\)](#), organochlorine pesticides, [bisphenol A](#) and some metals. The controversy around the effects of [endocrine disruptors](#) is perhaps best illustrated by research on [bisphenol A](#) (181, 182) whose estrogenic

activity was first reported in 1936. It was subsequently found to stabilize polycarbonates and resins and is now widely used in many products including food-can liners. There is a growing body of evidence related to the very low-dose effects of [bisphenol A](#) (66, 67, 183). The very low-dose effects of [endocrine disruptors](#) can not be predicted from high-dose studies, which contradicts the standard “dose makes the poison” rule of toxicology. Nontraditional dose-response curves are referred to as [nonmonotonic dose-response curves](#).

### 4.3.2 Fluoride

Fluoride is commonly added to municipal drinking water across the United States based on strong data that it reduces dental decay. This practice is supported by the US Centers for Disease Control (CDC) (184, 185). In addition to drinking water, fluoride is also present in a range of consumer products including toothpaste (1,000-1,500 parts per million or ppm), mouthwashes and fluoride supplements. The drinking water standards were established prior to the introduction of fluoride into many consumer products and the direct application of fluoride by dentists.

The central question, which is still unresolved, is what level of exposure results in harmful health effects to children. Children's small size means that, pound-for-pound of body weight, they receive a greater dose of fluoride than adults. The CDC estimates that up to 33% of children may have dental fluorosis because of the excessive intake of fluoride either through drinking water or through other sources -- an estimate which is supported by other studies (186). This concern has resulted in CDC issued a recommendation to limit fluoride exposure in children under eight years of age and to use fluoride-free water when preparing infant milk formula.

In addition, some recent studies suggest that excessive ingestion of fluoride lowers thyroid hormone levels, which is particularly critical for women with subclinical hypothyroidism. Decreased maternal thyroid levels are known to adversely affect fetal neurodevelopment (51). A study in China, for example, reported decreased child IQ levels associated with fluoride in drinking water. (Children in this study were also exposed to arsenic in drinking water, which may have confounded the results) (184, 185). The same year, in *Fluoride in Drinking Water: A Scientific Review of EPA's Standards*, a report produced by the National Academy of Sciences (NAS) (184), researchers analyzed the appropriateness of EPA's four ppm maximum contaminant level goal for fluoride in drinking water. The NAS was not directed to conduct a risk assessment of the effects of low-level fluoride exposure nor analyze other sources of exposure to fluoride. Referring to human and animal studies related to neurobehavioral effects, the NAS reports states “the consistency of the results appears significant enough to warrant additional research on the effects of fluoride on intelligence.” The primary question remains as to whether exposures to fluoride via multiple routes of exposure, from drinking water, food and dental-care products, may result in a high enough cumulative exposure to contribute to developmental effects.

While it is clear that fluoride has beneficial effects on dental health, emerging science suggests we need to further study the dose at which fluoridation may increase risks of neurodevelopment disorders, cancer and skeletal or dental fluorosis, particularly for sensitive individuals.

### 4.3.3 Food additives

Artificial or synthetic food colors and additives are ubiquitous in the food supply and have long been suspected of causing [conduct disorders](#). Their use has encouraged treatments such as the Feingold Diet (187, 188) in which many food additives are removed from the diet of individuals with [ADHD](#). Previous and recent carefully conducted double-blind human studies have confirmed that artificial food colorings such as sunset yellow, tartrazine, carmoisine and ponceau, as well as the preservative sodium benzoate, can cause [conduct disorders](#) (187-193). Recent studies using well-designed randomized, double-blind, placebo-controlled, crossover trials show that artificial food colors and additives cause increased

hyperactivity in three-year-old children (191). This has the potential to become a serious issue given the large number of children diagnosed with [ADHD](#).

## **5 Call for Further Research**

Further research into the links between [environmental agents](#) and [LDDs](#) is urgently needed, as is regulatory action in those cases in which the weight of evidence is sufficient now, to reduce exposures. Cumulative exposures (from different chemicals with similar modes of action) and aggregate exposures (from all sources of exposure to a chemical such as via dust, food, air, water), as well as direct exposure from items in the home environment and from food, must be taken into account in assessing risk and devising appropriate action.

### **5.1 Better assessment tools and procedures**

While some progress in testing methodology has been made, few chemicals have been evaluated using the Developmental Neurotoxicity Test (DNT) to date, although it has been determined as one of the most sensitive tests, in some cases showing adverse effects at lower levels of exposure than other tests. There is a growing demand to gather data on and screen compounds for effects on the nervous system. More sophisticated methodologies for testing humans and animals are also required, including testing procedures that address specific neurological disorders. New methods will be necessary to gather data as most high-volume chemicals have little or no developmental or neurotoxicity data.

### **5.2 In vitro (test-tube) screening of both new and old compounds**

For most chemicals there are very little data on the potential to cause learning or developmental disorders or even basic data on potential neurotoxicity. Basic research is needed to develop reliable and reproducible [in vitro](#) tests that accurately predict the potential for [neurotoxic](#) effects. Development of these tests will also help reduce the need for animal-based research studies. Relatively quick and inexpensive [in vitro](#) tests will need to be followed by more efficient, integrated neurobehavioral test methods if indicated (see the National Research Council report on toxicity testing (194)).

### **5.3 Chronic effects of hazardous chemicals**

While the acute high-dose effects of exposure to some chemicals are well documented, the learning and developmental effects of chronic, low-level exposure to pesticides, endocrine disruptors, flame retardants and other chemicals need further research.

### **5.4 Multiple exposures**

More research is needed on the consequences of exposure to multiple chemicals and cumulative exposures as children and adults are exposed to low levels of a variety of chemical agents throughout life. More data are needed on the interaction of the chemicals and their effects on development, and better risk-assessment procedures are needed to evaluate multiple exposures (195).

### **5.5 Low-level exposures**

Many animal and human studies have examined the effects of relatively high exposures. There is increasing evidence that low-level exposures to a variety of environmental toxicants (such as lead or tobacco) are associated with adverse consequences. In some cases the deficits are proportionately greater at lower levels. These new data indicates that studies should be designed specifically to examine the effects in representative samples, such as in the [National Children's Study](#).

## **5.6 Interactions with socioeconomic factors (environmental justice)**

Disadvantaged groups are at a higher risk for learning and developmental disorders. More research is needed to determine the mechanisms and relationship by which socioeconomic factors and stress interact with chemical exposures to produce learning and developmental disorders. Furthermore, we must develop tools and systemic models for prevention.

## **5.7 Effects of endocrine disruption on cognitive deficits**

Children and adults are exposed to a wide range of chemicals that affect the endocrine system and which can cause a range of learning and developmental disorders. Additional research is needed to evaluate the effects of low-level exposures and ascertain possible mechanisms of action and better characterize the adverse effects.

## **5.8 Interactions with genetics and identification of susceptible subpopulations**

There is increased need to identify subpopulations who, due to genetic contributors and susceptibility to developmental disorders, may be vulnerable to chemical exposure or other factors, including stress, that exacerbate the onset of these disorders.

## **5.9 World Health Organization research recommendations**

Research recommendations on evaluating children's risk to exposure to environmental chemicals are defined in a recent report by the World Health Organization (196).

# **6 Ethical and Policy Considerations**

## **6.1 Ethical considerations**

There is growing recognition that ethical, legal and social considerations play a crucial role in public- and child-health decision making that involves conflicts between individual, corporate, human rights and social-justice goals (197-201). Knowledge of the causes of learning and developmental disabilities implies an ethical duty and responsibility to act to protect children's health and well-being (122, 202). Accepting childhood exposure to contaminants that result in compromised learning and behavioral abilities violates the basic tenets of biomedical ethics. The principle of beneficence ("do good") requires that the benefits be maximized while the harm be minimized or eliminated. Respect for autonomy or personhood is violated when children are unnecessarily exposed to harmful substances. Respect of person also implies informed consent, and no child has given the informed consent for exposure to harmful chemicals. Finally, the principle of justice requires that burdens be shared equally, and because children are more vulnerable they endure a greater burden. In addition there are disparities related to socioeconomic status demonstrated by the increased burden of lead exposure in children of poverty (203). Perhaps America's first bioethicist Aldo Leopold said it best when he wrote in 1949: "A thing is right when it tends to preserve the integrity, stability, and beauty of the biotic community. It is wrong when it tends otherwise" (204). It is wrong to allow the exposure of children to environmental agents that cause learning and developmental disorders.

## **6.2 Policy considerations**

Recognition of the contribution of chemical contaminants to [LDDs](#) has increased substantially in recent years as new evidence has emerged both about the ability of neurotoxic chemicals to interfere with brain development and the susceptibility of the brain to chemicals (29). Given this established knowledge, it is clear that protecting children from [neurotoxic](#) environmental exposures from the earliest stages of fetal

development is an essential public health measure if we are to help prevent [LDDs](#) and create an environment in which children can reach and maintain their full potential.

There is a vast amount of information already available upon which to base sound policy decisions. As Garrett Hardin observed in 1968, many problems cannot be solved by technical solutions or additional research but only through responsible management of the problem (205). Our society is still contending with the effects of adding lead to paint and gasoline, even though its toxic effects were well documented at the time. Knowledge was sufficient, but management was not. To protect children, a precautionary approach is required that shifts the burden of responsibility to producers or manufacturers to demonstrate safety prior to potential exposure.

The researchers and reviewers for this statement are developing a companion document outlining specific policy recommendations based on the status of scientific knowledge outlined in this statement.

## **7 Conclusions**

The scientific evidence we have reviewed indicates environmental contaminants are an important cause of [learning and developmental disabilities](#). The proportion of environmentally induced [LDDs](#) is a question of profound human, scientific and public policy significance. Existing animal and human data suggest that a greater proportion is environmentally influenced than has yet been generally realized or than can be demonstrated with scientific certainty.

The consequences of [LDDs](#) are most significant for the affected individual but also have profound implications for the family, school system, local community and greater society. Despite some uncertainty, there is sufficient knowledge to take preventive action to reduce fetal and childhood exposures to environmental contaminants. Given the serious consequences of [LDDs](#), a precautionary approach is warranted to protect the most vulnerable of our society.

## 8 Resources: Children and Environmental Agents

American Association on Intellectual and  
Developmental Disabilities (AAIDD)  
[www.aaid.org/](http://www.aaid.org/)

American Pediatric Association  
[www.aap.org](http://www.aap.org)

Autism Society of America  
[www.autism-society.org](http://www.autism-society.org)

Center for Health, Environment and Justice  
[www.chej.org/](http://www.chej.org/)

Children's Environmental Health Network  
[www.cehn.org](http://www.cehn.org)

Collaborative on Health and the Environment  
[www.healthandenvironment.org](http://www.healthandenvironment.org)

Greater Boston Physicians for Social  
Responsibility  
<http://psr.igc.org/>

Healthy Child, Healthy World  
<http://healthychild.org/>

Healthy Schools Network, Inc.  
[www.healthyschools.org](http://www.healthyschools.org)

Institute for Children's Environmental Health  
[www.iceh.org](http://www.iceh.org)

Learning and Developmental Disabilities  
Initiative  
[www.iceh.org/LDDI.html](http://www.iceh.org/LDDI.html)

Learning Disabilities Association of America's  
Healthy Children Project  
[www.healthychildrenproject.org](http://www.healthychildrenproject.org)

Learning Disabilities Association of Canada  
(LDAC)  
[www.ldac-taac.ca](http://www.ldac-taac.ca)

Mt. Sinai Children's Environmental Health  
Center  
[www.cehcenter.org/](http://www.cehcenter.org/)

The National Association for the Dually  
Diagnosed  
[www.thenadd.org](http://www.thenadd.org)

National Institute for Environmental Health  
Science Centers for Children's Environmental  
Health & Disease Prevention Research  
[www.niehs.nih.gov/research/supported/centers/prevention/](http://www.niehs.nih.gov/research/supported/centers/prevention/)

Preventing Harm: A Resource and Action  
Center on Children and the Environment  
[www.preventingharm.org](http://www.preventingharm.org)

Toxipedia  
<http://toxipedia.org/>

University of Tennessee Youth Environment and  
Health Research Group  
<http://utyeah.utk.edu>

US Environmental Protection Agency Office of  
Children's Health Protection  
<http://yosemite.epa.gov/ochp/ochpweb.nsf/content/homepage.htm>

## 9 Glossary of Terms and Term Usage Specific to This Statement

(A number of the definitions were taken in whole or in part from the Vallombrosa Consensus Statement on Environmental Contaminants and Human Fertility Compromise October 2005.)

**Attention deficit hyperactivity disorder (ADHD)** – The principal characteristics of ADHD are inattention, hyperactivity and impulsivity. According to the most recent version of the Diagnostic and Statistical Manual of Mental Disorders ([DSM-IV-TR](#)) (206), there are three patterns of behavior that indicate ADHD. People with ADHD may show several signs of being consistently inattentive. They may have a pattern of being hyperactive and impulsive far more than others of their age. Or they may show both types of behavior. This means that there are three subtypes of ADHD recognized by professionals. These are the predominantly hyperactive-impulsive type (that does not show significant inattention); the predominantly inattentive type (that does not show significant hyperactive-impulsive behavior) sometimes called ADD – an outdated term for this entire disorder; and the combined type (that displays both inattentive and hyperactive-impulsive symptoms) (see also a document from the National Institute of Mental Health (13)).

**Autism spectrum disorders** – The Diagnostic and Statistical Manual of Mental Disorders [DSM-IV-TR](#) (206) includes autism as one of the five pervasive developmental disorders (PDD), more often referred to today as autism spectrum disorders (ASD). All these disorders are characterized by varying degrees of impairment in communication skills, social interactions and restricted, repetitive and stereotyped patterns of behavior (see also a document from the National Institute of Mental Health (16)).

**Bioaccumulation** – A process whereby contaminants taken up from the surrounding environment (air, food, water) are retained and concentrate in tissues at a rate faster than they can be broken down and excreted. With bioaccumulation, tissue levels of a contaminant become greater than surrounding environmental levels.

**Biomagnification** – A process whereby the concentration of contaminants increases up the food chain due to larger organisms ingesting smaller organisms containing contaminants. Humans and other predatory organisms accumulate the highest concentrations of contaminants.

**Biomarker** – A biological substance found in body fluids (blood, urine, breast milk) or tissues (fat) that can be measured and is associated with exposure to a contaminant. Biomarkers can help monitor exposure to contaminants and may help characterize individual susceptibilities to exposure. A biomarker of exposure is a measure of either the contaminant or a metabolite occurring shortly after exposure. A biomarker of effect is an enduring genetic change caused by a contaminant exposure that can be measured by changes in DNA or chromosome structures (such as [genetic mutations](#)). Biomarkers of effect are not necessarily specific to contaminant exposure. A biomarker of susceptibility is a gene or expression of a gene ([polymorphism](#)) that renders an individual more vulnerable to the adverse effects of contaminant exposure. For example, due to differences in enzymes some individuals may not be able to detoxify a contaminant as efficiently as others, resulting in higher levels of exposure and greater toxicity.

**Biomonitoring** – The assessment of exposure to contaminants by measuring [biomarkers](#) of exposure in body tissues or fluids (such as blood, urine, breast milk, amniotic fluid, hair, adipose tissue, bone). Biomonitoring can be used to monitor not only exposures in populations but also changes in levels of contaminants over time.

**Bisphenol A** – A common chemical compound that forms the building block of polycarbonate plastics and epoxy resins. Bisphenol A is used in polycarbonate plastic in food containers, water bottles, baby

bottles, compact disk cases, eyeglass lenses, the lining of food cans and as a dental sealant. It binds with nuclear and extracellular estrogen receptors.

**Conduct disorders** – Refers to behavior characterized by hostility and aggression.

**Developmental delays** – Refers to development which is delayed compared with age peers. Developmental delays can be associated with varied or negative effects in multiple areas such as sensory, motor, language, social, reading and emotional areas. These are also referred to as neurodevelopmental delays.

**Dioxins** – A class of hundreds of related [persistent](#) chemicals, some of which are known to be highly toxic, that result from industrial combustion/incineration processes; burning of household trash or fuels such as wood, coal and oil; chlorine bleaching of pulp/paper; and some types of chemical manufacturing. Cigarette smoke also contains dioxins.

**DSM-IV-TR** – The Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR) (206) is a handbook for mental health professionals that lists categories of mental disorders and the criteria for diagnosing them according to the American Psychiatric Association. It is used worldwide by clinicians and researchers as well as insurance companies, pharmaceutical companies and policymakers.

**Dyslexia** – A specific learning disability that manifests primarily as a difficulty with written language, particularly with reading and spelling. Dyslexia is the result of a neurological difference but is not an intellectual disability. Most people with dyslexia have average or above-average intelligence.

**Endocrine disruptors** – Environmental compounds that interfere with the normal function of endogenous hormones (those produced by an organism). Endocrine disruptors can stimulate or block the actions of hormones or can interfere with their metabolism. Endocrine disruptors continue to be discovered but have been recognized to include a diverse range of chemicals including pesticides, plasticizers, flame retardants, industrial byproducts, pharmaceuticals and plant-derived compounds.

**Endpoints** – In a scientific study, the outcome that is being measured. In the field of environmental health, this could be a [biomarker](#), toxic effect, disease outcome or other measure anticipated to differ between exposed and unexposed populations.

**Environmental agents** – These include natural or synthetic chemicals, heavy metals (such as lead, mercury, cadmium), and naturally occurring compounds such as plant-derived estrogens.

**Environmental factors** – In this document, “environmental factors” refers to a broader range of possible environmental influences than the environmental agents listed above. Environmental factors include pharmaceutical use, stress, other chemical agents or physical conditions that adversely affect learning or development.

**Epigenetics** – The molecular phenomena that regulate gene expression without alterations to the DNA sequence.

**Genetic mutation** – A change in the nucleotide sequence of a DNA molecule. Genetic mutations are a kind of [genetic polymorphism](#). Genetic mutation refers to changes in DNA sequence which are not present in most individuals of a species and either have been associated with disease (or risk of disease) or have resulted from damage inflicted by external agents (such as viruses or radiation).

**Genetic Polymorphism** – A difference in DNA sequence among individuals, groups or populations (for example, a genetic polymorphism might increase susceptibility to Fetal Alcohol Spectrum Disorder). Genetic polymorphisms may be the result of chance processes, may be inherited or may be induced by external agents (such as viruses or radiation), which is then referred to as a genetic mutation.

**In vitro testing** – Generally refers to experiments done in a test tube, outside a living organism. *In vitro* systems typically use cell culture techniques for initial assessment of harmful effects.

**Learning and developmental disabilities (LDDs)** – Conditions resulting from interference of normal brain development and function that adversely affect an individual’s performance. [Learning and developmental disabilities](#) include but are not limited to deficits in learning and memory, reduced IQ, [attention deficit hyperactivity disorder](#), [autism spectrum disorder](#), [conduct disorders](#) and [developmental delays](#), but do not consider effects of the peripheral nervous system.

**Mental retardation** – According to the American Association on Intellectual and Developmental Disabilities (AAIDD), mental retardation is a disability characterized by significant limitations both in intellectual functioning and in adaptive behavior as expressed in conceptual, social and practical adaptive skills (see [www.aaid.org/Policies/faq\\_mental\\_retardation.shtml](http://www.aaid.org/Policies/faq_mental_retardation.shtml)).

**Mixtures** – In this context, “mixtures” indicates the effects of two or more contaminants in which the outcome of exposure is different from their separate effects. The interaction could be additive (a sum of individual effects), subtractive (one substance is stimulatory and another inhibitory), or multiplicative (the effect is greater than the sum of individual effects).

**Morphometric** – Generally refers to procedures that count or quantitatively assess the number of specific cell types.

**National Health and Nutrition Examination Survey (NHANES)** – A program designed to assess the health and nutritional status of adults and children in the United States. The survey is unique in that it combines interviews and physical examinations. NHANES is a major program of the National Center for Health Statistics (NCHS), which is part of the Centers for Disease Control and Prevention (CDC), US Public Health Service, and has the responsibility for producing vital and health statistics for the nation ([www.cdc.gov/nchs/nhanes.htm](http://www.cdc.gov/nchs/nhanes.htm)).

**National Children’s Study** – Led by a consortium of US government agencies (<http://nationalchildrenstudy.gov>), this study aims to examine the effects of [environmental factors](#) on the health and development of more than 100,000 children from before birth to age 21.

**Neuropathology/neuropathological** – The microscopic study of cells of the brain or nervous system.

**Neurotoxic / neurotoxicity** – An adverse change in the chemistry, structure or function of the nervous system, during development or at maturity, following exposure to a chemical or physical agent.

**Neurotoxicant** – A chemical or physical agent that produces [neurotoxicity](#).

**Nonmonotonic dose-response curve (NMDR)** – A traditional dose-response curve in toxicology assumes that the response to exposure will increase with increasing dose. This is known as a monotonic curve, one in which the slope of the dose-response curve does not change from positive to negative or vice versa. In a nonmonotonic dose-response curve, the slope of the dose-response curve changes sign as the level of exposure increases. Some NMDR curves are shaped like a U, while others are shaped like an inverted U. NMDR curves are important from a public-health perspective because in dose-response

curves that are nonmonotonic, low-dose effects cannot be predicted from high-dose testing. The traditional assumption that higher doses cause greater harm (“the dose makes the poison”) is used in standard risk-assessment studies to identify the level of a chemical exposure beneath which contamination should cause no effect. This old assumption may be true for many chemicals and for many classic health effects, but it can be misleading for exposures that have a nonmonotonic dose-response curve.

**Octyl/nonyl phenols** – Chemicals that belong to a broader class of compounds known as alkylphenol ethoxylates (APEs). APEs are high-volume chemicals that have been used for more than 40 years as detergents, emulsifiers and wetting and dispersing agents. These chemicals are used as ingredients in spermicides, cosmetics and detergents and as inert ingredients in pesticides. Some are [endocrine disruptors](#). Several are noted contaminants in aquatic environments.

**Perfluorinated compounds (PFCs)** – [Persistent, bioaccumulative](#) chemicals found in a wide array of products including stain-resistant coatings for carpets and clothing (Gore-Tex), nonstick cookware (Teflon), and insecticides. Widespread contamination of human tissues has been documented, with some of the highest levels found in US populations. PFCs have been linked to neuroendocrine and reproductive effects.

**Persistence** – This refers to the stability of a contaminant in the environment. Persistent contaminants are characterized by their ability to resist natural degradation so that they build up in the environment with time. Persistent contaminants often are transported globally on currents of wind or water.

**Phthalates** – Chemicals added to personal-care products to enhance penetration and hold scent/color and used as plasticizers in rigid plastics such as polyvinyl chloride (PVC) to create flexibility. Phthalates are found in numerous and diverse consumer products including vinyl flooring, plastic shower curtains, cosmetics and fragrances, shampoos and lotions, toys, pharmaceutical and herbal pill coatings – and in hospital equipment including intravenous bags and tubing.

**Polybrominated diphenyl ethers (PBDEs)** – [Persistent, bioaccumulative](#) chemicals added to electronics, upholstery foam, textiles and numerous other materials to make them more flame-resistant. PBDEs have a chemical structure very similar to [PCBs](#) and have been rapidly accumulating in wildlife and human tissues.

**Polychlorinated biphenyls (PCBs)** – There are 209 individual chlorinated PCB compounds (known as congeners) that are [persistent](#) and [bioaccumulative](#). Manufacture was banned in the US in the late 1970s, although PCBs are still found in some products, and widespread environmental contamination still exists. PCBs were used in hundreds of commercial and industrial applications, including as lubricants, plasticizers, insulators for electrical applications, caulking and paint. Health effects of PCB exposure include acne-like skin conditions in adults and neurobehavioral and immunological changes in children. PCBs cause cancer in animals (see ATSDR fact sheet [www.atsdr.cdc.gov/tfacts17.html](http://www.atsdr.cdc.gov/tfacts17.html)).

**Polymorphism** – see genetic polymorphism above.

**Potentiated or potentiate** – To enhance or increase the effect of a drug or chemical, which results in an increased response or undesired action or effect.

## **10 Acknowledgments**

This document would not exist but for the drive and vision of Elise Miller, MEd, executive director of the Institute for Children’s Environmental Health (ICEH) and national coordinator of the Collaborative on Health and the Environment’s Learning and Developmental Disabilities Initiative, as well as the persistence and expertise of Steven G. Gilbert, PhD, DABT. A special acknowledgment goes to Nancy Snow, MS, research and communications manager at ICEH, who consistently provided thoughtful editing and invaluable organizational comments and always responded throughout numerous drafts. Sean Foley of Toxipedia and the Institute for Neurotoxicology and Neurological Disorders was instrumental helping with early research and editorial comments, and A. T. Birmingham-Young of Precise Words Editing Services provided final copyediting. We are particularly indebted to Bernie Weiss, Ted Schettler, Philip Landrigan, Bruce Lanphear, Martha Herbert, and Barbara McElgunn for their in-depth comments on various aspects of the document throughout the editing process. And finally, thank you to the entire review team for their formal and informal suggestions, encouragement and support in conceiving and developing this document.

## 11 References

1. Dick FD. Parkinson's disease and pesticide exposures. *Br Med Bull* 2006;79-80:219-31.
2. Dick FD, Seaton A, Haites N, Semple SE, Dick S, Prescott GJ, et al. Environmental risk factors for Parkinson's disease and parkinsonism: the Geoparkinson study. *Occup Environ Med* 2007.
3. Richardson JR, Caudle WM, Wang M, Dean ED, Pennell KD, Miller GW. Developmental exposure to the pesticide dieldrin alters the dopamine system and increases neurotoxicity in an animal model of Parkinson's disease. *Faseb J* 2006;20(10):1695-7.
4. Singh C, Ahmad I, Kumar A. Pesticides and metals induced Parkinson's disease: involvement of free radicals and oxidative stress. *Cell Mol Biol (Noisy-le-grand)* 2007;53(5):19-28.
5. Schmidt CW. Environmental connections: a deeper look into mental illness. *Environ Health Perspect* 2007;115(8):404-410.
6. Gilbert SG, Grant-Webster KS. Neurobehavioral effects of developmental methylmercury exposure. *Environ Health Perspect* 1995;103 Suppl 6:135-42.
7. Boyle CA, Decoufle P, Yeargin-Allsopp M. Prevalence and health impact of developmental disabilities in US children. *Pediatrics* 1994;93(3):399-403.
8. Perrin JM, Bloom SR, Gortmaker SL. The increase of childhood chronic conditions in the United States. *Jama* 2007;297(24):2755-9.
9. Biasini FJ, Grupe L, Huffman L, Bray NW. Mental retardation: a symptom and a syndrome. In: Netherton SD, Holmes D, Walker CE, editors. *Child and Adolescent Psychological Disorders: A Comprehensive Textbook (Oxford Textbooks in Clinical Psychology, Vol 4)*. USA: Oxford University Press; 1999.
10. Davidson P, Myers C, Weiss B, editors. *Neurotoxicity and Developmental Disabilities*: Academic Press; 2006.
11. Larson SA, Lakin KC, Anderson L, Kwak N, Lee JH, Anderson D. Prevalence of mental retardation and developmental disabilities: estimates from the 1994/1995 National Health Interview Survey Disability Supplements. *Am J Ment Retard* 2001;106(3):231-52.
12. Schroeder SR. Mental retardation and developmental disabilities influenced by environmental neurotoxic insults. *Environ Health Perspect* 2000;108 Suppl 3:395-9.
13. NIMH. Attention Deficit Hyperactivity Disorder. In: Health NIOm, editor.: National Institute of Mental Health; 2006.
14. Goldman LS, Genel M, Bezman RJ, Slanetz PJ. Diagnosis and treatment of attention-deficit/hyperactivity disorder in children and adolescents. Council on Scientific Affairs, American Medical Association. *Jama* 1998;279(14):1100-7.
15. Froehlich T, Lanphear B, Epstein J, Barbaresi W, Katusic S, Kahn R. Prevalence, recognition and treatment of attention-deficit, hyperactivity disorder in a national sample of US children. *Arch Ped Adolesc Med* 2007;in press.
16. NIMH. Autism Spectrum Disorders (Pervasive Developmental Disorders). In: Health NIOm, editor.: National Institute of Mental Health; 2004.
17. Branum AM, Schoendorf KC. Changing patterns of low birthweight and preterm birth in the United States, 1981-98. *Paediatr Perinat Epidemiol* 2002;16(1):8-15.
18. Braun JM, Kahn RS, Froehlich T, Auinger P, Lanphear BP. Exposures to environmental toxicants and attention deficit hyperactivity disorder in US children. *Environ Health Perspect* 2006;114(12):1904-9.

19. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM. Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. *JAMA* 2004;291(23):2847-50.
20. Landrigan PJ, Schechter CB, Lipton JM, Fahs MC, Schwartz J. Environmental pollutants and disease in American children: estimates of morbidity, mortality, and costs for lead poisoning, asthma, cancer, and developmental disabilities. *Environ Health Perspect* 2002;110(7):721-8.
21. Davies K. Economic costs of childhood diseases and disabilities attributable to environmental contaminants in Washington State, USA. *EcoHealth* 2006;3(2):86-94.
22. Chambers JG, Parrish T, Harr JJ. What are we spending on special education services in the United States, 1999-2000?, Advance Report #1: American Institutes for Research; 2002 March 2002.
23. Gilbert SG, Weiss B. A rationale for lowering the blood lead action level from 10 to 2 microg/dL. *Neurotoxicology* 2006;27(5):693-701.
24. Clarkson T. Methylmercury and fish consumption: weighing the risks. *Can Med Assoc J* 1998;158:1465-1466.
25. Clarkson TW. The three modern faces of mercury. *Environ Health Perspect* 2002;110 Suppl 1:11-23.
26. Debes F, Budtz-Jorgensen E, Weihe P, White RF, Grandjean P. Impact of prenatal methylmercury exposure on neurobehavioral function at age 14 years. *Neurotoxicol Teratol* 2006;28(5):536-47.
27. Grandjean P, Weihe P, White RF, Debes F, Araki S, Yokoyama K, et al. Cognitive deficit in 7-year-old children with prenatal exposure to methylmercury. *Neurotoxicol Teratol* 1997;19(6):417-28.
28. Grandjean P, White RF. Effects of methylmercury exposure on neurodevelopment. *Jama* 1999;281(10):896; author reply 897.
29. Grandjean P, Landrigan PJ. Developmental neurotoxicity of industrial chemicals. *Lancet* 2006;368(9553):2167-78.
30. Antoniadis A, Gilbert SG, Gagnon M. Neurotoxicants: environmental contributors to disability in children. *The ASHA Leader - American Speech-Language-Hearing Association* 2006;11(13):6-7, 38-39.
31. Miller E, Snow N. Safeguarding our children at home: reducing exposures to toxic chemicals and heavy metals. *Zero to Three* 2005;November:26-32.
32. Orzech D. Chemical kids — environmental toxins and child development. *Social Work Today* 2007;7(2):37-40.
33. Perera FP, Illman SM, Kinney PL, Whyatt RM, Kelvin EA, Shepard P, et al. The challenge of preventing environmentally related disease in young children: community-based research in New York City. *Environ Health Perspect* 2002;110(2):197-204.
34. Cory-Slechta DA, Virgolini MB, Thiruchelvam M, Weston DD, Bauter MR. Maternal stress modulates the effects of developmental lead exposure. *Environ Health Perspect* 2004;112(6):717-30.
35. van den Hazel P, Zuurbier M, Babisch W, Bartonova A, Bistrup ML, Bolte G, et al. Today's epidemics in children: possible relations to environmental pollution and suggested preventive measures. *Acta Paediatr Suppl* 2006;95(453):18-25.
36. Weiss B, Bellinger DC. Social ecology of children's vulnerability to environmental pollutants. *Environ Health Perspect* 2006;114(10):1479-85.

37. Linnet KM, Dalsgaard S, Obel C, Wisborg K, Henriksen TB, Rodriguez A, et al. Maternal lifestyle factors in pregnancy risk of attention deficit hyperactivity disorder and associated behaviors: review of the current evidence. *Am J Psychiatry* 2003;160(6):1028-40.
38. Grandjean P, Bellinger D, Bergman A, Cordier S, Davey-Smith G, Eskenazi B, et al. The faroes statement: human health effects of developmental exposure to chemicals in our environment. *Basic & Clinical Pharmacology & Toxicology* 2007;0(0):1-3.
39. Landrigan P, Nordberg M, Lucchini R, Nordberg G, Grandjean P, Iregren A, et al. The Declaration of Brescia on prevention of the neurotoxicity of metals. *Am J Ind Med* 2007;50(10):709-11.
40. Schettler T, Stein J, Reich F, Valenti M. In *Harm's Way: Toxic Threats to Child Development*. Cambridge, MA: Greater Boston Physicians for Social Responsibility; 2000.
41. Sokol RJ, Delaney-Black V, Nordstrom B. Fetal alcohol spectrum disorder. *JAMA* 2003;290(22):2996-9.
42. Adams J, Barone S, Jr., LaMantia A, Philen R, Rice DC, Spear L, et al. Workshop to identify critical windows of exposure for children's health: neurobehavioral work group summary. *Environ Health Perspect* 2000;108 Suppl 3:535-44.
43. Rice D, Barone S, Jr. Critical periods of vulnerability for the developing nervous system: evidence from humans and animal models. *Environ Health Perspect* 2000;108 Suppl 3:511-33.
44. Rodier PM. Vulnerable periods and processes during central nervous system development. *Environ Health Perspect* 1994;102 Suppl 2:121-4.
45. Rodier PM. Developing brain as a target of toxicity. *Environ Health Perspect* 1995;103 Suppl 6:73-6.
46. Clarkson TW. Mercury: major issues in environmental health. *Environ Health Perspect* 1992;100:31-38.
47. Cory-Slechta DA. Studying toxicants as single chemicals: does this strategy adequately identify neurotoxic risk? *Neurotoxicology* 2005;26(4):491-510.
48. Okey AB, Franc MA, Moffat ID, Tijet N, Boutros PC, Korkalainen M, et al. Toxicological implications of polymorphisms in receptors for xenobiotic chemicals: the case of the aryl hydrocarbon receptor. *Toxicol Appl Pharmacol* 2005;207(2 Suppl):43-51.
49. Palanza P, Morellini F, Parmigiani S, vom Saal FS. Prenatal exposure to endocrine disrupting chemicals: effects on behavioral development. *Neurosci Biobehav Rev* 1999;23(7):1011-27.
50. Schantz SL, Widholm JJ. Cognitive effects of endocrine-disrupting chemicals in animals. *Environ Health Perspect* 2001;109(12):1197-206.
51. LaFranchi SH, Haddow JE, Hollowell JG. Is thyroid inadequacy during gestation a risk factor for adverse pregnancy and developmental outcomes? *Thyroid* 2005;15(1):60-71.
52. Oerbeck B, Sundet K, Kase BF, Heyerdahl S. Congenital hypothyroidism: influence of disease severity and L-thyroxine treatment on intellectual, motor, and school-associated outcomes in young adults. *Pediatrics* 2003;112(4):923-30.
53. Selva KA, Harper A, Downs A, Blasco PA, Lafranchi SH. Neurodevelopmental outcomes in congenital hypothyroidism: comparison of initial T4 dose and time to reach target T4 and TSH. *J Pediatr* 2005;147(6):775-80.
54. Warren KR, Li TK. Genetic polymorphisms: impact on the risk of fetal alcohol spectrum disorders. *Birth Defects Res A Clin Mol Teratol* 2005;73(4):195-203.

55. Furlong CE, Cole TB, Jarvik GP, Pettan-Brewer C, Geiss GK, Richter RJ, et al. Role of paraoxonase (PON1) status in pesticide sensitivity: genetic and temporal determinants. *Neurotoxicology* 2005;26(4):651-9.
56. Furlong CE, Holland N, Richter RJ, Bradman A, Ho A, Eskenazi B. PON1 status of farmworker mothers and children as a predictor of organophosphate sensitivity. *Pharmacogenet Genomics* 2006;16(3):183-90.
57. Holland N, Furlong C, Bastaki M, Richter R, Bradman A, Huen K, et al. Paraoxonase polymorphisms, haplotypes, and enzyme activity in Latino mothers and newborns. *Environ Health Perspect* 2006;114(7):985-91.
58. Kelada SN, Shelton E, Kaufmann RB, Houry MJ. Delta-aminolevulinic acid dehydratase genotype and lead toxicity: a HuGE review. *Am J Epidemiol* 2001;154(1):1-13.
59. Montenegro MF, Barbosa F, Jr., Sandrim VC, Gerlach RF, Tanus-Santos JE. A polymorphism in the delta-aminolevulinic acid dehydratase gene modifies plasma/whole blood lead ratio. *Arch Toxicol* 2006;80(7):394-8.
60. Anway MD, Skinner MK. Epigenetic transgenerational actions of endocrine disruptors. *Endocrinology* 2006;147(6 Suppl):S43-9.
61. Jones PA, Takai D. The role of DNA methylation in mammalian epigenetics. *Science* 2001;293(5532):1068-70.
62. Skinner MK, Anway MD. Epigenetic transgenerational actions of vinclozolin on the development of disease and cancer. *Crit Rev Oncog* 2007;13(1):75-82.
63. Anway MD, Leathers C, Skinner MK. Endocrine disruptor vinclozolin induced epigenetic transgenerational adult-onset disease. *Endocrinology* 2006;147(12):5515-23.
64. Anway MD, Memon MA, Uzumcu M, Skinner MK. Transgenerational effect of the endocrine disruptor vinclozolin on male spermatogenesis. *J Androl* 2006;27(6):868-79.
65. Newbold RR, Padilla-Banks E, Jefferson WN. Adverse effects of the model environmental estrogen diethylstilbestrol are transmitted to subsequent generations. *Endocrinology* 2006;147(6 Suppl):S11-7.
66. vom Saal FS, Hughes C. An extensive new literature concerning low-dose effects of bisphenol A shows the need for a new risk assessment. *Environ Health Perspect* 2005;113(8):926-33.
67. vom Saal FS, Nagel SC, Timms BG, Welshons WV. Implications for human health of the extensive bisphenol A literature showing adverse effects at low doses: a response to attempts to mislead the public. *Toxicology* 2005;212(2-3):244-52, author reply 253-4.
68. Deutch B, Pedersen HS, Asmund G, Hansen JC. Contaminants, diet, plasma fatty acids and smoking in Greenland 1999-2005. *Sci Total Environ* 2007;372(2-3):486-96.
69. Huwe JK, Larsen GL. Polychlorinated dioxins, furans, and biphenyls, and polybrominated diphenyl ethers in a US meat market basket and estimates of dietary intake. *Environ Sci Technol* 2005;39(15):5606-11.
70. NHANES. National Health and Nutrition Examination Survey. Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention National Center for Health Statistics; 2007.
71. Rodier PM, Ingram JL, Tisdale B, Croog VJ. Linking etiologies in humans and animal models: studies of autism. *Reprod Toxicol* 1997;11(2-3):417-22.

72. Hallene KL, Oby E, Lee BJ, Santaguida S, Bassanini S, Cipolla M, et al. Prenatal exposure to thalidomide, altered vasculogenesis, and CNS malformations. *Neuroscience* 2006;142(1):267-83.
73. Sadamatsu M, Kanai H, Xu X, Liu Y, Kato N. Review of animal models for autism: implication of thyroid hormone. *Congenit Anom (Kyoto)* 2006;46(1):1-9.
74. Miyazaki K, Narita N, Narita M. Maternal administration of thalidomide or valproic acid causes abnormal serotonergic neurons in the offspring: implication for pathogenesis of autism. *Int J Dev Neurosci* 2005;23(2-3):287-97.
75. Herbert MR, Russo JP, Yang S, Roohi J, Blaxill M, Kahler SG, et al. Autism and environmental genomics. *Neurotoxicology* 2006;27(5):671-84.
76. Muhle R, Trentacoste SV, Rapin I. The genetics of autism. *Pediatrics* 2004;113(5):e472-86.
77. Mahaffey KR. Nutrition and lead: strategies for public health. *Environ Health Perspect* 1995;103 Suppl 6:191-6.
78. Golub MS, Hogrefe CE, Germann SL, Tran TT, Beard JL, Crinella FM, et al. Neurobehavioral evaluation of rhesus monkey infants fed cow's milk formula, soy formula, or soy formula with added manganese. *Neurotoxicol Teratol* 2005;27(4):615-27.
79. Davidsson L, Cederblad A, Lonnerdal B, Sandstrom B. The effect of individual dietary components on manganese absorption in humans. *Am J Clin Nutr* 1991;54(6):1065-70.
80. Hu H, Kotha S, Brennan T. The role of nutrition in mitigating environmental insults: policy and ethical issues. *Environ Health Perspect* 1995;103 Suppl 6:185-90.
81. Lau K, McLean WG, Williams DP, Howard CV. Synergistic interactions between commonly used food additives in a developmental neurotoxicity test. *Toxicol Sci* 2006;90(1):178-87.
82. Tiffany-Castiglioni E, editor. *In Vitro Neurotoxicology: Principles and Challenges (Methods in Pharmacology and Toxicology)* Humana Press; 2004.
83. Bolon B, Garman R, Jensen K, Krinke G, Stuart B. A 'best practices' approach to neuropathologic assessment in developmental neurotoxicity testing--for today. *Toxicol Pathol* 2006;34(3):296-313.
84. Moser VC. The functional observational battery in adult and developing rats. *Neurotoxicology* 2000;21(6):989-96.
85. EPA. Health Effects Test Guidelines. OPPTS 870.6300 Developmental Neurotoxicity Study. In: US Environmental Protection Agency; 1998.
86. OECD. Draft guidance document on reproductive toxicity testing and assessment, Series on Testing and Assessment No. 43. Environment Directorate, Organisation for Economic Cooperation and Development, November 10, 2004. Available: <http://www.oecd.org/dataoecd/38/46/34030071.pdf>. 2004.
87. Claudio L, Kwa WC, Russell AL, Wallinga D. Testing methods for developmental neurotoxicity of environmental chemicals. *Toxicol Appl Pharmacol* 2000;164(1):1-14.
88. Cory-Slechta DA, Crofton KM, Foran JA, Ross JF, Sheets LP, Weiss B, et al. Methods to identify and characterize developmental neurotoxicity for human health risk assessment. I: behavioral effects. *Environ Health Perspect* 2001;109 Suppl 1:79-91.
89. Dorman DC, Allen SL, Byczkowski JZ, Claudio L, Fisher JE, Jr., Fisher JW, et al. Methods to identify and characterize developmental neurotoxicity for human health risk assessment. III: pharmacokinetic and pharmacodynamic considerations. *Environ Health Perspect* 2001;109 Suppl 1:101-11.

90. Garman RH, Fix AS, Jortner BS, Jensen KF, Hardisty JF, Claudio L, et al. Methods to identify and characterize developmental neurotoxicity for human health risk assessment. II: neuropathology. *Environ Health Perspect* 2001;109 Suppl 1:93-100.
91. Milesion BE, Ferenc SA. Methods to identify and characterize developmental neurotoxicity for human health risk assessment: overview. *Environ Health Perspect* 2001;109 Suppl 1:77-8.
92. Burbacher TM, Grant KS. Methods for studying nonhuman primates in neurobehavioral toxicology and teratology. *Neurotoxicol Teratol* 2000;22(4):475-86.
93. Moser VC, Simmons JE, Gennings C. Neurotoxicological interactions of a five-pesticide mixture in preweanling rats. *Toxicol Sci* 2006;92(1):235-45.
94. Ehman KD, Moser VC. Evaluation of cognitive function in weanling rats: a review of methods suitable for chemical screening. *Neurotoxicol Teratol* 2006;28(1):144-61.
95. Cory-Slechta DA. Lead-induced impairments in complex cognitive function: offerings from experimental studies. *Child Neuropsychol* 2003;9(1):54-75.
96. Burbacher TM, Grant KS, Mayfield DB, Gilbert SG, Rice DC. Prenatal methylmercury exposure affects spatial vision in adult monkeys. *Toxicol Appl Pharmacol* 2005;208(1):21-8.
97. Rice DC, Gilbert SG. Early chronic low-level methylmercury poisoning in monkeys impairs spatial vision. *Science* 1982;216(4547):759-61.
98. Rice DC, Gilbert SG. Exposure to methylmercury from birth to adulthood impairs high-frequency hearing in monkeys. *Toxicol Appl Pharmacol* 1992;115(1):6-10.
99. Rice DC, Gilbert SG. Effects of developmental methylmercury exposure or lifetime lead exposure on vibration sensitivity function in monkeys. *Toxicol Appl Pharmacol* 1995;134(1):161-9.
100. Rice DC. Behavioral deficit (delayed matching to sample) in monkeys exposed from birth to low levels of lead. *Toxicol Appl Pharmacol* 1984;75(2):337-45.
101. Rice DC, Gilbert SG. Sensitive periods for lead-induced behavioral impairment (nonspatial discrimination reversal) in monkeys. *Toxicol Appl Pharmacol* 1990;102(1):101-9.
102. Rice DC. Neurotoxicity of lead, methylmercury, and PCBs in relation to the Great Lakes. *Environ Health Perspect* 1995;103 Suppl 9:71-87.
103. Grandjean P, Weihe P, Burse VW, Needham LL, Storr-Hansen E, Heinzow B, et al. Neurobehavioral deficits associated with PCB in 7-year-old children prenatally exposed to seafood neurotoxicants. *Neurotoxicol Teratol* 2001;23(4):305-17.
104. Anger WK. Neurobehavioural tests and systems to assess neurotoxic exposures in the workplace and community. *Occup Environ Med* 2003;60(7):531-8, 474.
105. Farahat TM, Abdelrasoul GM, Amr MM, Shebl MM, Farahat FM, Anger WK. Neurobehavioural effects among workers occupationally exposed to organophosphorous pesticides. *Occup Environ Med* 2003;60(4):279-86.
106. Kamel F, Rowland AS, Park LP, Anger WK, Baird DD, Gladen BC, et al. Neurobehavioral performance and work experience in Florida farmworkers. *Environ Health Perspect* 2003;111(14):1765-72.
107. McCauley LA, Anger WK, Keifer M, Langley R, Robson MG, Rohlman D. Studying health outcomes in farmworker populations exposed to pesticides. *Environ Health Perspect* 2006;114(6):953-60.

108. Binder LM, Storzbach D, Anger WK, Campbell KA, Rohlman DS, Center HR. Subjective cognitive complaints, affective distress, and objective cognitive performance in Persian Gulf War veterans. *Arch Clin Neuropsychol* 1999;14(6):531-6.
109. Binder LM, Storzbach D, Campbell KA, Rohlman DS, Anger WK. Neurobehavioral deficits associated with chronic fatigue syndrome in veterans with Gulf War unexplained illnesses. *J Int Neuropsychol Soc* 2001;7(7):835-9.
110. Anger WK, Liang YX, Nell V, Kang SK, Cole D, Bazylewicz-Walczak B, et al. Lessons learned -15 years of the WHO-NCTB: a review. *Neurotoxicology* 2000;21(5):837-46.
111. Rohlman DS, Anger WK, Tamulinas A, Phillips J, Bailey SR, McCauley L. Development of a neurobehavioral battery for children exposed to neurotoxic chemicals. *Neurotoxicology* 2001;22(5):657-65.
112. Lanphear BP, Hornung R, Khoury J, Yolton K, Baghurst P, Bellinger DC, et al. Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environ Health Perspect* 2005;113(7):894-9.
113. Needleman HL, Bellinger D. The health effects of low level exposure to lead. *Annu Rev Public Health* 1991;12:111-140.
114. Myers GJ, Davidson PW, Cox C, Shamlaye C, Cernichiari E, Clarkson TW. Twenty-seven years studying the human neurotoxicity of methylmercury exposure. *Environ Res* 2000;83(3):275-85.
115. Schettler T. Toxic threats to neurologic development of children. *Environ Health Perspect* 2001;109 Suppl 6:813-6.
116. Warren KR, Calhoun FJ, May PA, Viljoen DL, Li TK, Tanaka H, et al. Fetal alcohol syndrome: an international perspective. *Alcohol Clin Exp Res* 2001;25(5 Suppl ISBRA):202S-206S.
117. Streissguth AP, Barr HM, Sampson PD. Moderate prenatal alcohol exposure: effects on child IQ and learning problems at age 7 1/2 years. *Alcohol Clin Exp Res* 1990;14(5):662-9.
118. Streissguth AP, Barr HM, Sampson PD, Bookstein FL. Prenatal alcohol and offspring development: the first fourteen years. *Drug Alcohol Depend* 1994;36(2):89-99.
119. Canfield RL, Henderson CR, Jr., Cory-Slechta DA, Cox C, Jusko TA, Lanphear BP. Intellectual impairment in children with blood lead concentrations below 10 microg per deciliter. *N Engl J Med* 2003;348(16):1517-26.
120. Miranda ML, Kim D, Galeano MA, Paul CJ, Hull AP, Morgan SP. The Relationship between early childhood blood lead levels and performance on end-of-grade tests. *Environ Health Perspect* 2007;115(8):1242-7.
121. Bellinger DC, Bellinger AM. Childhood lead poisoning: the torturous path from science to policy. *J Clin Invest* 2006;116(4):853-7.
122. Gilbert SG. Ethical, legal, and social issues: our children's future. *Neurotoxicology* 2005;26(4):521-530.
123. Davidson PW, Myers GJ, Weiss B, Shamlaye CF, Cox C. Prenatal methylmercury exposure from fish consumption and child development: a review of evidence and perspectives from the Seychelles Child Development Study. *Neurotoxicology* 2006;27(6):1106-9.
124. Davidson PW, Myers GJ, Cox C, Wilding GE, Shamlaye CF, Huang LS, et al. Methylmercury and neurodevelopment: longitudinal analysis of the Seychelles child development cohort. *Neurotoxicol Teratol* 2006;28(5):529-35.

125. Trasande L, Landrigan PJ, Schechter C. Public health and economic consequences of methylmercury toxicity to the developing brain. *Environ Health Perspect* 2005;113(5):590-6.
126. Goldman LR, Farland WH. Methylmercury risks. *Science* 1998;279(5351):640-1; author reply 641.
127. Grandjean P, White RF, Nielsen A, Cleary D, de Oliveira Santos EC. Methylmercury neurotoxicity in Amazonian children downstream from gold mining. *Environ Health Perspect* 1999;107(7):587-91.
128. ATSDR. Toxicological Profile For Polychlorinated Biphenyls (PCBs): Agency for Toxic Substances and Disease Registry. Available at <http://www.atsdr.cdc.gov/toxprofiles/tp17.pdf>. 2000.
129. Jacobson JL, Jacobson SW. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *N Engl J Med* 1996;335(11):783-9.
130. Jacobson JL, Jacobson SW. Dose-response in perinatal exposure to polychlorinated biphenyls (PCBs): the Michigan and North Carolina cohort studies. *Toxicol Ind Health* 1996;12(3-4):435-45.
131. Nakajima S, Saijo Y, Kato S, Sasaki S, Uno A, Kanagami N, et al. Effects of prenatal exposure to polychlorinated biphenyls and dioxins on mental and motor development in Japanese children at 6 months of age. *Environ Health Perspect* 2006;114(5):773-8.
132. Vreugdenhil HJ, Van Zanten GA, Brocaar MP, Mulder PG, Weisglas-Kuperus N. Prenatal exposure to polychlorinated biphenyls and breastfeeding: opposing effects on auditory P300 latencies in 9-year-old Dutch children. *Dev Med Child Neurol* 2004;46(6):398-405.
133. Schantz SL, Gasior DM, Polverejan E, McCaffrey RJ, Sweeney AM, Humphrey HE, et al. Impairments of memory and learning in older adults exposed to polychlorinated biphenyls via consumption of Great Lakes fish. *Environ Health Perspect* 2001;109(6):605-11.
134. Schantz SL, Widholm JJ, Rice DC. Effects of PCB exposure on neuropsychological function in children. *Environ Health Perspect* 2003;111(3):357-576.
135. Costa LG, Giordano G. Developmental neurotoxicity of polybrominated diphenyl ether (PBDE) flame retardants. *Neurotoxicology* 2007.
136. Rice DC, Reeve EA, Herlihy A, Zoeller RT, Thompson WD, Markowski VP. Developmental delays and locomotor activity in the C57BL6/J mouse following neonatal exposure to the fully-brominated PBDE, decabromodiphenyl ether. *Neurotoxicol Teratol* 2007;29(4):511-20.
137. Bowler RM, Nakagawa S, Drezgic M, Roels HA, Park RM, Diamond E, et al. Sequelae of fume exposure in confined space welding: a neurological and neuropsychological case series. *Neurotoxicology* 2007;28(2):298-311.
138. Bowler RM, Roels HA, Nakagawa S, Drezgic M, Diamond E, Park R, et al. Dose-effect relationships between manganese exposure and neurological, neuropsychological and pulmonary function in confined space bridge welders. *Occup Environ Med* 2007;64(3):167-77.
139. ATSDR. Toxicological Profile For Manganese: Agency for Toxic Substances and Disease Registry. Available at <http://www.atsdr.cdc.gov/toxprofiles/tp17.pdf>. 2000.
140. Bouchard M, Laforest F, Vandelac L, Bellinger D, Mergler D. Hair manganese and hyperactive behaviors: pilot study of school-age children exposed through tap water. *Environ Health Perspect* 2007;115(1):122-7.
141. Wasserman GA, Liu X, Parvez F, Ahsan H, Factor-Litvak P, Kline J, et al. Water arsenic exposure and intellectual function in 6-year-old children in Araihaazar, Bangladesh. *Environ Health Perspect* 2007;115(2):285-9.

142. Woolf AD, Wright RO, Amarasiriwardena C, Bellinger DC. A child with chronic manganese exposure from drinking water. *Environ Health Perspect* 2002;110(6):613-6.
143. Wright RO, Amarasiriwardena C, Woolf AD, Jim R, Bellinger DC. Neuropsychological correlates of hair arsenic, manganese, and cadmium levels in school-age children residing near a hazardous waste site. *Neurotoxicology* 2006;27(2):210-6.
144. Pal PK, Samii A, Calne DB. Manganese neurotoxicity: a review of clinical features, imaging and pathology. *Neurotoxicology* 1999;20(2-3):227-38.
145. Mergler D, Baldwin M, Belanger S, Larribe F, Beuter A, Bowler R, et al. Manganese neurotoxicity, a continuum of dysfunction: results from a community based study. *Neurotoxicology* 1999;20(2-3):327-42.
146. Wasserman GA, Liu X, Parvez F, Ahsan H, Levy D, Factor-Litvak P, et al. Water manganese exposure and children's intellectual function in Araihasar, Bangladesh. *Environ Health Perspect* 2006;114(1):124-9.
147. Ahsan H, Chen Y, Parvez F, Zablotska L, Argos M, Hussain I, et al. Arsenic exposure from drinking water and risk of premalignant skin lesions in Bangladesh: baseline results from the Health Effects of Arsenic Longitudinal Study. *Am J Epidemiol* 2006;163(12):1138-48.
148. Wasserman GA, Liu X, Parvez F, Ahsan H, Factor-Litvak P, van Geen A, et al. Water arsenic exposure and children's intellectual function in Araihasar, Bangladesh. *Environ Health Perspect* 2004;112(13):1329-33.
149. Hafeman DM, Ahsan H, Louis ED, Siddique AB, Slavkovich V, Cheng Z, et al. Association between arsenic exposure and a measure of subclinical sensory neuropathy in Bangladesh. *J Occup Environ Med* 2005;47(8):778-84.
150. Golub MS, Macintosh MS, Baumrind N. Developmental and reproductive toxicity of inorganic arsenic: animal studies and human concerns. *J Toxicol Environ Health B Crit Rev* 1998;1(3):199-241.
151. ATSDR. Toxicological Profile For Arsenic: Agency for Toxic Substances and Disease Registry. Available at <http://www.atsdr.cdc.gov/toxprofiles/tp17.pdf>. 2005.
152. White RF, Proctor SP. Solvents and neurotoxicity. *Lancet* 1997;349(9060):1239-43.
153. Eskenazi B, Gaylord L, Bracken MB, Brown D. In utero exposure to organic solvents and human neurodevelopment. *Dev Med Child Neurol* 1988;30(4):492-501.
154. Laslo-Baker D, Barrera M, Knittel-Keren D, Kozer E, Wolpin J, Khattak S, et al. Child neurodevelopmental outcome and maternal occupational exposure to solvents. *Arch Pediatr Adolesc Med* 2004;158(10):956-61.
155. Till C, Westall CA, Koren G, Nulman I, Rovet JF. Vision abnormalities in young children exposed prenatally to organic solvents. *Neurotoxicology* 2005;26(4):599-613.
156. Till C, Westall CA, Rovet JF, Koren G. Effects of maternal occupational exposure to organic solvents on offspring visual functioning: a prospective controlled study. *Teratology* 2001;64(3):134-41.
157. Arnold GL, Kirby RS, Langendoerfer S, Wilkins-Haug L. Toluene embryopathy: clinical delineation and developmental follow-up. *Pediatrics* 1994;93(2):216-20.
158. Hersh JH, Podruch PE, Rogers G, Weisskopf B. Toluene embryopathy. *J Pediatr* 1985;106(6):922-7.
159. Choi H, Jedrychowski W, Spengler J, Camann DE, Whyatt RM, Rauh V, et al. International studies of prenatal exposure to polycyclic aromatic hydrocarbons and fetal growth. *Environ Health Perspect* 2006;114(11):1744-50.

160. Perera FP, Rauh V, Whyatt RM, Tsai WY, Tang D, Diaz D, et al. Effect of prenatal exposure to airborne polycyclic aromatic hydrocarbons on neurodevelopment in the first 3 years of life among inner-city children. *Environ Health Perspect* 2006;114(8):1287-92.
161. Guillette EA, Meza MM, Aquilar MG, Soto AD, Garcia IE. An anthropological approach to the evaluation of preschool children exposed to pesticides in Mexico. *Environ Health Perspect* 1998;106(6):347-53.
162. Rohlman DS, Arcury TA, Quandt SA, Lasarev M, Rothlein J, Travers R, et al. Neurobehavioral performance in preschool children from agricultural and non-agricultural communities in Oregon and North Carolina. *Neurotoxicology* 2005;26(4):589-98.
163. Berkowitz GS, Wetmur JG, Birman-Deych E, Obel J, Lapinski RH, Godbold JH, et al. In utero pesticide exposure, maternal paraoxonase activity, and head circumference. *Environ Health Perspect* 2004;112(3):388-91.
164. Grandjean P, Harari R, Barr DB, Debes F. Pesticide exposure and stunting as independent predictors of neurobehavioral deficits in Ecuadorian school children. *Pediatrics* 2006;117(3):e546-56.
165. Ruckart PZ, Kakolewski K, Bove FJ, Kaye WE. Long-term neurobehavioral health effects of methyl parathion exposure in children in Mississippi and Ohio. *Environ Health Perspect* 2004;112(1):46-51.
166. Whyatt RM, Garfinkel R, Hoepner LA, Holmes D, Borjas M, Williams MK, et al. Within- and between-home variability in indoor-air insecticide levels during pregnancy among an inner-city cohort from New York City. *Environ Health Perspect* 2007;115(3):383-9.
167. Whyatt RM, Rauh V, Barr DB, Camann DE, Andrews HF, Garfinkel R, et al. Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ Health Perspect* 2004;112(10):1125-32.
168. Young JG, Eskenazi B, Gladstone EA, Bradman A, Pedersen L, Johnson C, et al. Association between in utero organophosphate pesticide exposure and abnormal reflexes in neonates. *Neurotoxicology* 2005;26(2):199-209.
169. Handal AJ, Lozoff B, Breilh J, Harlow SD. Effect of community of residence on neurobehavioral development in infants and young children in a flower-growing region of Ecuador. *Environ Health Perspect* 2007;115(1):128-33.
170. Handal AJ, Lozoff B, Breilh J, Harlow SD. Neurobehavioral development in children with potential exposure to pesticides. *Epidemiology* 2007;18(3):312-20.
171. Handal AJ, Lozoff B, Breilh J, Harlow SD. Sociodemographic and nutritional correlates of neurobehavioral development: a study of young children in a rural region of Ecuador. *Rev Panam Salud Publica* 2007;21(5):292-300.
172. Rauh VA, Garfinkel R, Perera FP, Andrews HF, Hoepner L, Barr DB, et al. Impact of prenatal chlorpyrifos exposure on neurodevelopment in the first 3 years of life among inner-city children. *Pediatrics* 2006;118(6):e1845-59.
173. Orlebeke JF, Knol DL, Verhulst FC. Child behavior problems increased by maternal smoking during pregnancy. *Arch Environ Health* 1999;54(1):15-9.
174. Thapar A, Fowler T, Rice F, Scourfield J, van den Bree M, Thomas H, et al. Maternal smoking during pregnancy and attention deficit hyperactivity disorder symptoms in offspring. *Am J Psychiatry* 2003;160(11):1985-9.
175. Wakschlag LS, Pickett KE, Cook E, Jr., Benowitz NL, Leventhal BL. Maternal smoking during pregnancy and severe antisocial behavior in offspring: a review. *Am J Public Health* 2002;92(6):966-74.

176. Miller T, Rauh VA, Glied SA, Hattis D, Rundle A, Andrews H, et al. The economic impact of early life environmental tobacco smoke exposure: early intervention for developmental delay. *Environ Health Perspect* 2006;114(10):1585-8.
177. Gatzke-Kopp LM, Beauchaine TP. Direct and Passive Prenatal Nicotine Exposure and the Development of Externalizing Psychopathology. *Child Psychiatry Hum Dev* 2007.
178. Wakschlag LS, Pickett KE, Leventhal BL. The association between maternal cigarette smoking and psychiatric diseases or criminal outcome in the offspring. *Reprod Toxicol* 2000;14(6):579-80.
179. Brennan PA, Grekin ER, Mortensen EL, Mednick SA. Relationship of maternal smoking during pregnancy with criminal arrest and hospitalization for substance abuse in male and female adult offspring. *Am J Psychiatry* 2002;159(1):48-54.
180. MMWR. Smoking During Pregnancy - United States, 1990-2002. In: Center for Disease Control and Prevention. Available at <http://www.cdc.gov/MMWR/preview/mmwrhtml/mm5339a1.htm>; 2004. p. 911-915.
181. Gross L. The toxic origins of disease. *PLoS Biol* 2007;5(7):e193.
182. Kuehn BM. Expert panels weigh bisphenol-A risks. *JAMA* 2007;298(13):1499-503.
183. vom Saal FS, Cooke PS, Buchanan DL, Palanza P, Thayer KA, Nagel SC, et al. A physiologically based approach to the study of bisphenol A and other estrogenic chemicals on the size of reproductive organs, daily sperm production, and behavior. *Toxicol Ind Health* 1998;14(1-2):239-60.
184. NRC. Fluoride in Drinking Water: A Scientific Review of EPA's Standards. In: National Research Council of the National Academies. Available at [http://books.nap.edu/openbook.php?record\\_id=11571&page=R1](http://books.nap.edu/openbook.php?record_id=11571&page=R1); 2006.
185. Wang SX, Wang ZH, Cheng XT, Li J, Sang ZP, Zhang XD, et al. Arsenic and fluoride exposure in drinking water: children's IQ and growth in Shanyin county, Shanxi province, China. *Environ Health Perspect* 2007;115(4):643-7.
186. Erdal S, Buchanan SN. A quantitative look at fluorosis, fluoride exposure, and intake in children using a health risk assessment approach. *Environ Health Perspect* 2005;113(1):111-7.
187. Weiss B, Williams JH, Margen S, Abrams B, Caan B, Citron LJ, et al. Behavioral responses to artificial food colors. *Science* 1980;207(4438):1487-9.
188. Schab DW, Trinh NH. Do artificial food colors promote hyperactivity in children with hyperactive syndromes? A meta-analysis of double-blind placebo-controlled trials. *J Dev Behav Pediatr* 2004;25(6):423-34.
189. Bateman B, Warner JO, Hutchinson E, Dean T, Rowlandson P, Gant C, et al. The effects of a double blind, placebo controlled, artificial food colourings and benzoate preservative challenge on hyperactivity in a general population sample of preschool children. In: *Arch Dis Child*; 2004. p. 506-11.
190. Boris M, Mandel FS. Foods and additives are common causes of the attention deficit hyperactive disorder in children. *Ann Allergy* 1994;72(5):462-8.
191. McCann D, Barrett A, Cooper A, Crumpler D, Dalen L, Grimshaw K, et al. Food additives and hyperactive behaviour in 3-year-old and 8/9-year-old children in the community: a randomised, double-blinded, placebo-controlled trial. *Lancet* 2007;370(9598):1560-7.
192. Rowe KS, Briggs DR. Food additives and behaviour. *Med J Aust* 1994;161(10):581-2.
193. Rowe KS, Rowe KJ. Synthetic food coloring and behavior: a dose response effect in a double-blind, placebo-controlled, repeated-measures study. *J Pediatr* 1994;125(5 Pt 1):691-8.

194. NRC. Toxicity Testing in the 21st Century: A Vision and a Strategy Committee on Toxicity Testing and Assessment of Environmental Agents: National Research Council of the National Academies. Available at <http://www.nap.edu/catalog/11970.html>; 2007.
195. Boekelheide K. Mixed Messages. *Toxicological Sciences* 2007;99(1):1-2.
196. WHO. Principles for evaluating health risks in children associated with exposure to chemicals (Environmental health criteria ; 237): World Health Organization; 2006.
197. Lee C. Environmental justice: building a unified vision of health and the environment. *Environ Health Perspect* 2002;110 Suppl 2:141-4.
198. Schettler TH. Human rights: necessary? sufficient? diversionary? *Int J Occup Environ Health* 2003;9(1):69-73.
199. Callahan D, Jennings B. Ethics and public health: forging a strong relationship. *Am J Public Health* 2002;92(2):169-76.
200. Kass NE. An ethics framework for public health. *Am J Public Health* 2001;91(11):1776-82.
201. Smith C. Human rights, environment, and individual action. *Int J Occup Environ Health* 2003;9(1):1-3.
202. Weiss B. Ethics assessment as an adjunct to risk assessment in the evaluation of developmental neurotoxicants. *Environ Health Perspect* 2001;109 Suppl 6:905-8.
203. Bernard SM, McGeehin MA. Prevalence of blood lead levels  $\geq 5$  microg/dL among US children 1 to 5 years of age and socioeconomic and demographic factors associated with blood of lead levels 5 to 10 microg/dL, Third National Health and Nutrition Examination Survey, 1988-1994. *Pediatrics* 2003;112(6 Pt 1):1308-13.
204. Leopold A. *A Sand County Almanac*: Oxford University Press, USA; 2 edition (December 31, 1968); 1949.
205. Hardin G. The tragedy of the commons. *Science* 1968;162(859):1243-8.
206. APA. *The Diagnostic and Statistical Manual of Mental Disorders. DSM-IV-TR*. 4th ed. Washington, DC: American Psychiatric Association; 2000.

**The following scientists, researchers and health professionals affirm that they each agree with content of the LDDI Scientific Consensus Statement on Environmental Agents Associated with Neurodevelopmental Disorders:**

**Anastasia Antoniadis, MPH**, Early Intervention, Devon, Pennsylvania

**David K. Becker, MD, MPH**, University of California at San Francisco

**David C. Bellinger, PhD, MSc**, Harvard Medical School /Children's Hospital Boston, Massachusetts

**JoAnn Rupert Behm, MS, RN**, Learning Disabilities Association of California, Novato

**Karen R. Bowman, MN, RN**, Karen Bowman & Associates, Inc., Seattle, Washington

**Asa Bradman, PhD, MS**, University of California, Berkeley

**Thomas Burbacher PhD**, University of Washington, Seattle

**Janice E. Camp, MSN, MSPH, CIH**, Field Research and Consultation Group, University of Washington, Seattle

**David O. Carpenter, MD**, University at Albany, State University of New York

**Lilian Corra, MD**, International Society of Doctors for the Environment, Argentina

**Christine A. Curran, PhD**, University of Cincinnati, Ohio

**Lee M. Dorigan, MES, RS**, Public Health – Seattle & King County, Washington

**Lori L. Driscoll, PhD**, Colorado College, Colorado Springs, Colorado

**Steven G. Gilbert, PhD, DABT**, Institute of Neurotoxicology & Neurological Disorders, Seattle, Washington

**Lynn R. Goldman, MD, MPH**, Johns Hopkins University, Baltimore, Maryland

**Philippe Grandjean, MD, DMSc**, Harvard University, Boston, Massachusetts

**Kimberly S. Grant, PhD**, University of Washington, Seattle

**Martha Reed Herbert, MD, PhD**, Massachusetts General Hospital, Boston

**Bonnie J. Kaplan, PhD**, University of Calgary, Alberta

**Catherine J. Karr, MD, PhD**, University of Washington, Seattle

**Wenceslao A. Kiat, MD**, St. Luke's Medical Center, Quezon City, Philippines

**Janna Koppe, MD, PhD**, Ecobaby Foundation, Loenersloot, The Netherlands

**Stephen R. Krival, MS**, University of Washington, Seattle

**Philip J. Landrigan, MD, MSc**, Mount Sinai School of Medicine, New York, New York

**Bruce Lanphear, MD, MPH**, Cincinnati Children's Hospital Medical Center, Ohio

**Eric V. Larsson, PhD, LP, BCBA**, Lovaas Institute Midwest, Minneapolis, Minnesota

**Susan Makris, MS**, US Environmental Protection Agency, Washington, DC

**Kathleen Matula, PhD**, University of Texas, San Antonio

**Barbara McElgunn, RN**, Learning Disabilities Association of Canada, Ottawa

**Kelly Marie McVeary, PhD**, Georgetown University, Washington, DC

**Donna Mergler, PhD**, University of Québec at Montréal

**John Peterson Myers, PhD**, Environmental Health Sciences, Charlottesville, Virginia

**Virginia C. Moser, PhD, DABT**, Research Triangle Park, North Carolina

**Gina Muckle, PhD**, Laval University, Sainte-Foy, Québec

**M. Christopher Newland, PhD**, Auburn University, Alabama

**MaryAnn O'Hara, MD, MPH, MSt**, University of Washington, Seattle

**Rebecca L. Parant, MS, RN**, Bradley Hospital, East Providence, Rhode Island

**Lynne Parsons Heilbrun, MPH**, University of Texas Health Science Center at San Antonio

**Joan Patterson, PhD**, University of Minnesota, Minneapolis

**Isaac Pessah, PhD**, University of California, Davis

**Janet Primomo, PhD, RN**, University of Washington, Tacoma

**Edward P. Riley, PhD**, San Diego State University, California

**John Roberts, MS, MEd, PE**, Engineering Plus, Sammamish, Washington

**Lawrence D. Rosen, MD**, Old Tappan Pediatrics and Hackensack University Medical Center, New Jersey

**Sheela Sathyanarayana, MD, MPH**, University of Washington, Seattle

**Ted Schettler, MD, MPH**, Science and Environmental Health Network, Newburyport, Massachusetts

**Margaret Shield, PhD**, Local Hazardous Waste Management Program in King County, Seattle, Washington

**Gregg Stanwood, PhD**, Vanderbilt University, Nashville, Tennessee

**Jill Stein, MD**, Greater Boston Physicians for Social Responsibility, Massachusetts

**Sandra Steingraber, PhD**, Ithaca College, New York

**Julia Helen Tracy, MS**, University of Washington, Seattle

**Tatjana T. Walker, RD, CDE**, University of Texas Health Science Center at San Antonio

**Charles V. Vorhees, PhD**, University of Cincinnati, Ohio

**David Wallinga, MD, MPA**, Healthy Legacy, Minneapolis, Minnesota

**Bernard Weiss, PhD**, University of Rochester, New York

**Christine Ziebold, MD, PhD, MPH**, University of Iowa, Iowa City

June 3, 2008